

Geometric invariant theory approach to the determination of ground states of D-wave condensates in isotropic space

Yu.M. Gufan¹, Al.V. Popov¹, G. Sartori^{2,4}, V. Talamini^{3,4}, G. Valente^{2,4} and E.B. Vinberg⁵

¹Rostov State University, Rostov on Don, Russia

²Dipartimento di Fisica, Università di Padova, Italy

³Dipartimento di Ingegneria Civile, Università di Udine, Italy

⁴INFN, Sezione di Padova, Italy

⁵ Moscow State University, Moscow, Russia

A complete and rigorous determination of the possible ground states for D-wave pairing Bose condensates is presented, using a geometrical invariant theory approach to the problem. The order parameter is argued to be a vector, transforming according to a ten dimensional real representation of the group $G = \mathbf{O}_3 \otimes \mathbf{U}_1 \times \langle T \rangle$. We determine the equalities and inequalities defining the orbit space of this linear group and its symmetry strata, which are in a one-to-one correspondence with the possible distinct phases of the system. We find 15 allowed phases (besides the unbroken one), with different symmetries, that we thoroughly determine. The group-subgroup relations between bordering phases are pointed out. The perturbative sixth degree corrections to the minimum of a fourth degree polynomial G -invariant free energy, calculated by Mermin, are also determined.

PACS 74.20.De, 02.20.-a, 64.60.-i, MSC 14L24

I. INTRODUCTION

Superfluidity and superconductivity are justified on the basis of the macroscopic condensation of Bose quasi-particles. The classical Bardeen, Cooper and Schrieffer theory for superconductivity dates 1957. Soon after, a BCS-type transition was proposed for the Fermi system ^3He by Anderson and Morel [1]. Cooper pair formation was thought to occur in an $L \neq 0$ state, to take into account the hard core nature of ^3He atom interaction. The superfluid phases were actually observed [2], and the nature of p-wave pairing is now well established for ^3He . The theory of $L \neq 0$ -superfluids is relevant for “unconventional” superconductivity too. The high temperature superconducting (HTS) oxides are anomalous in their non-Fermi liquid normal state properties and in many cases share with heavy fermion superconductors unconventional d-wave pairing behaviour [3,4]. In fact, experiments probing the phase and the nodes of the gap show a sign reversal of the order parameter, compatible with the d-wave scenario. Nevertheless, other experiments on the same HTS compounds point in the direction of a significant s-wave component. In the presence of external magnetic fields, defects and interface phenomena there are also evidences of a mixed pairing symmetry (see *e.g.*, the Introduction of [5] and references therein). Such a controversial experimental situation is reflected in the fact that the underlying microscopic mechanism inducing superconductivity in HTS materials is still unclear and is one of today’s major challenges [6,7].

Such a situation has motivated the efforts at studying the macroscopic properties of unconventional superconductors through the Landau theory of phase transitions [8] and the identification of the order parameter symmetry has become a preliminary task in the construction of viable models describing the attractive nature of the pairing interaction.

Direct experimental evidence of an order parameter unconventional structure lies on multiple phase transitions. The heavy fermion compounds $\text{U}_{1-x}\text{Th}_x\text{Be}_{13}$ display four different superconducting phases in the T - x phase diagram and UPt_3 displays three superconducting phases in a T - H phase diagram [3]. Furthermore, from power-law temperature behaviour of thermodynamic and transport properties (*e.g.* specific heat, magnetic penetration depth), a non-trivial node structure for the gap function may be inferred, compatible with high L -pairing. That is the case also for HTS oxides, for which, at present, clear proofs are lacking of the existence of more than one superconducting state.

Actually, it must be pointed out that, unlike superfluid ^3He , which is an isotropic fermion system, Bloch electrons in a superconductor crystal lattice exhibit, in general, a reduced finite symmetry; in addition, the existence of imperfections, even in the cleanest samples, can partly destroy the gap node structure at low energy. So, the very fact that the experimental observation cannot at present completely unravel the node structure of the gap function, reinforces the necessity of classifying all the possible symmetry breaking schemes.

Moreover, the role of Fermi isotropic space beyond a zero-th order approximation in phenomenological theories, is still supported by some recent studies: In HTS oxides, low symmetrical crystal fields (tetragonal and orthorhombic), as well as spontaneous strain, have weak influence on the temperature of the superconducting phase transition [9] or on the penetration depth [10]. So, even if the spectrum of the interactions which are responsible for the pairing

(whatever their origins may be) must be anisotropic in the crystal, it is worth analyzing the possibility that they act in a way not directly dependent on the crystal structure (like in isotropic space).

The possible ground states of a high L superfluid has been the object of intense investigations during the 60's and the 70's. From the solution of the state equations, Anderson and Morel [1] and Mermin [11] identified five different phases, through the minimization of a 4-th degree Landau potential. Schakel and Bias [12] analyzed the problem using only group theoretical arguments. Capel and Schakel [13], taking advantage of the results in [12], computed the possible ground states of condensates driven by p-waves. They also investigated the consequences of lowering the residual symmetry by means of strong spin-orbit interactions. Since, in this case, the Cooper pair is in a $J = 2$ state ($S = 1, L = 1$), the order parameter is represented by a traceless symmetric tensor. The same holds true for an ($S = 0, L = 2$) state order parameter, so their results may be directly applied to D-wave pairing. According to the authors of ref. [13], eleven different phases are allowed. Their analysis, however, is questionable, since time reversal symmetry cannot be neglected in the theory of condensate states and the use of a 4-th degree polynomial Landau free energy strongly limits the number of phases that would be allowed by the symmetry of the system.

Our aim is to give a definitive answer to the classification of possible symmetry breaking patterns in D-wave pairing Bose condensate, in the framework of the Landau theory of phase transitions. In this paper, we shall mainly be concerned with the mathematical aspects of the problem, that will be solved making use of the geometrical invariant theory approach, proposed in ref. [14]. The strategy is to exploit a set of basic invariant polynomials of the symmetry group of the system, as fundamental variables in the description of the phase space of the system and in the minimization procedure of the free energy [15]. In the realization of this program, we shall meet some substantial computational difficulties, that will be overcome by means of innovative procedures.

II. THE GEOMETRICAL INVARIANT THEORY APPROACH TO SPONTANEOUS SYMMETRY BREAKING

Let us briefly recall some basic elements of the geometrical invariant theory approach, or orbit space approach (see [16] and references therein) to the determination of possible patterns of spontaneous symmetry breaking.

Let $x \in \mathbf{R}^n$ be a vector order parameter, transforming linearly and orthogonally under the compact real symmetry group G , and let $\Phi(\alpha; x)$ be the G -invariant free energy, expressed in terms, also, of state variables α . The points $x_0(\alpha)$, where the function $\phi_\alpha(x) = \Phi(\alpha; x)$ takes on its absolute minimum, determine the stable phase of the system, whose residual symmetry is defined by the isotropy subgroup of G at x_0 , G_{x_0} . Owing to its G -invariance, the free energy is a constant along each G -orbit, so, each of its stationary points is degenerate along the G -orbit through it. Since the isotropy subgroups of G at points of the same orbit are conjugate in G , only the conjugate class, $[G_{x_0}]$, of G_{x_0} in G , i.e. the *orbit-type* (or *symmetry*) of the orbit through x_0 , is physically relevant.

The set of all G -orbits, endowed with the quotient topology and differentiable structure, forms the *orbit space*, $\mathbf{R}^n/G$, of G and the subset of all the G -orbits with the same orbit-type forms a *stratum* of $\mathbf{R}^n/G$. Phase transitions take place when, by varying the values of the α 's, the point $x_0(\alpha)$ is shifted to an orbit lying on a different stratum.

If $\Phi(\alpha; x)$ is a sufficiently general function of the α 's, by varying these parameters, the point $x_0(\alpha)$ can be shifted to any stratum of $\mathbf{R}^n/G$. So, *the strata are in a one-to-one correspondence with the symmetry phases allowed by the G -invariance of the free energy*. On the contrary, extra restrictions on the form of the free energy function, not coming from G -symmetry requirements (e.g., the assumption that the free energy is a polynomial of low degree), can limit the number of allowed phases for the system in its ground state.

Being constant along each G -orbit, the free energy may be conveniently thought of as a function defined in the orbit space of G . This fact can be formalized using some basic results of invariant theory. In fact, the G -invariant polynomial functions separate the G -orbits, meaning that, for any two distinct G -orbits, there is at least a polynomial G -invariant function assuming different values on them. Moreover, every G -invariant polynomial can be built as a real polynomial function of a *finite* set, $\{p_1(x), \dots, p_q(x)\}$, of basic polynomial invariants (*integrity basis of the ring of G -invariant polynomials*). Thus, the elements of an integrity basis can be conveniently used as coordinates of the orbit space points. They need not, for general compact groups, be algebraically independent, but can, and will, be chosen to be homogeneous polynomials in x . The number of algebraically independent elements in a *minimal* set of basic polynomial invariants is $n - \nu$, where ν is the dimension of a generic (principal) orbit of G . Information on the number and degrees of a minimal set of basic invariants, and the degrees of the algebraic relations (*syzygies*) among them, can be inferred from the Mölién function of G .

Let us call q_0 the number of independent elements of the set $\{p\}$. The range of the *orbit map*, $x \mapsto p(x) = (p_1(x), \dots, p_q(x)) \in \mathbf{R}^q$, yields a realization of the orbit space of the linear group G , as a connected semi-algebraic surface, i.e. a subset of $\mathbf{R}^q$, determined by algebraic equations and inequalities. The orbit space of G will, therefore, be identified with a closed and connected region, $\hat{S}$, of a q_0 -dimensional algebraic surface, delimited by lower dimensional

semi-algebraic surfaces.

If an integrity basis has been determined, the equations and inequalities defining the orbit space of a compact group can be obtained from a simple recipe. It has been shown, in fact, that the orbit space of a reductive linear group can be identified with the semialgebraic variety formed by the points $p \in \mathbf{R}^q$, satisfying the following conditions i) and ii) [14,17]:

- i) p lies on the surface, Z , defined by the syzygies;
- ii) the $q \times q$ matrix $\hat{P}(p)$, defined by the relations

$$\hat{P}_{ab}(p(x)) = \sum_{j=1}^n \partial_j p_a(x) \partial_j p_b(x), \quad \forall x \in \mathbf{R}^n \quad (1)$$

is positive semi-definite and has rank q_0 at p .

The minimum of $\Phi(\alpha; x)$ can be computed as a *constrained* minimum of the function, $\hat{\Phi}(\alpha; p)$, $p \in \hat{S}$, defined by

$$\hat{\Phi}(\alpha; p(x)) = \Phi(\alpha; x), \quad \forall x \in \mathbf{R}^n. \quad (2)$$

It has also been shown [14] that the state equation

$$\partial \Phi(\alpha; x) / \partial x_j = 0, \quad j = 1, \dots, n,$$

determining the extremal points of $\Phi(\alpha; x)$, is equivalent to the following equation in orbit space:

$$\sum_{b=1}^q \hat{P}_{ab}(p) \partial_b \hat{\Phi}(\alpha; p) = 0, \quad a = 1, \dots, q, \quad p \in \hat{S}. \quad (3)$$

Like all semi-algebraic sets, the orbit space of G presents a natural *stratification*. It can, in fact, be considered as the disjoint union of a *finite number* of semi-algebraic subsets of decreasing dimensions (*geometrical strata*), each geometrical stratum being in the border of a higher dimensional one, but for the highest dimensional stratum, which is unique (*principal stratum*). The *geometrical strata* are the connected components of the *symmetry strata*. The symmetries of two bordering strata are related by a group-subgroup relation and the orbit-type of the lower dimensional stratum is larger.

In order to determine the minimum of $\hat{\Phi}(\alpha; p)$, $p \in \hat{S}$, as a constrained minimum, one needs the relations defining the geometrical strata of $\mathbf{R}^n/G$. These can be obtained from positivity and rank conditions on the matrix $\hat{P}(p)$ and from the syzygies, as already stated.

The geometrical and symmetry strata are determined in the following way. Let $\widehat{W}^{(d)}$ denote the (generally non connected) d dimensional algebraic subset of Z , defined by the relation

$$\widehat{W}^{(d)} = \{p \in Z \mid \text{rank}(\hat{P}(p)) = d, \hat{P}(p) \geq 0\}, \quad (4)$$

then, the geometrical d dimensional strata are the connected components of $\widehat{W}^{(d)}$ and each d dimensional symmetry stratum, $\hat{S}^{(d,\alpha)}$, $\alpha = 1, \dots$, is the union of all the geometrical d dimensional strata, $\widehat{W}^{(d,\alpha,r)}$, $r = 1, \dots, k_\alpha$, with the same orbit-type, $[G^{(d,r)}]$. A representative in $[G^{(d,r)}]$ can be obtained as the isotropy subgroup at an arbitrarily chosen point $x^{(d,r)}$ of the image in $\mathbf{R}^n$, $W^{(d,r)}$, of the geometric stratum $\widehat{W}^{(d,r)}$. The point $x^{(d,r)}$ can be obtained as a solution of the equation¹ $p(x) = p^{(d,r)}$, where $p^{(d,r)}$ is an arbitrarily chosen point in $\widehat{W}^{(d,r)}$.

In the following, we shall classify and characterize all the allowed symmetry phases and possible phase transitions between contiguous phases, for D -wave phase condensates in isotropic space. In particular, in Section III we shall identify the linear symmetry group, G , of these systems and a minimal set of basic polynomial invariants of G . In Section IV, we shall determine the geometrical features of the orbit space of G , *i.e.*, its stratification (including

¹For $p^{(0)} \in \hat{S}$, all the solutions, x , of the equation $\tilde{p}(x) = \tilde{p}^{(0)}$ form a unique G -orbit. The equation has no solutions for $p^{(0)} \notin \hat{S}$.

connection properties and bordering relations of the strata) and the orbit-types of its strata. In section V, using an innovative method, we shall provide relatively simple expressions for the equalities and inequalities determining each stratum. These are essential for establishing the connection properties of the strata and make much easier the determination of the minima of a specific free energy function. A simple example of the determination of the minima of a general 4-th degree G -invariant polynomial and their stability against 6-th order perturbations, will be given in Section V.

III. SYMMETRY OF THE ALLOWED D-WAVE CONDENSATE STATES IN ISOTROPIC SPACE

The formation of D-wave condensate ground states breaks the symmetry of the isotropic 3-dimensional space, which corresponds to the group $\mathbf{O}_3 \otimes \mathbf{U}_1 \times \langle \mathcal{T} \rangle$, where $\mathbf{O}_3$ is the complete rotation group, $\mathbf{U}_1$ is the gauge group and $\langle \mathcal{T} \rangle$ is the group² generated by the time reversal operator $\mathcal{T}$.

The symmetry of the allowed D-wave condensate ground states is defined by the relative values of the complex coefficients in the decomposition of the gap-function, Δ , in terms of spherical harmonics with $L = 2$:

$$\Delta(\theta, \phi) = \sum_{m=-2}^2 D_m Y_2^m(\theta, \phi).$$

The set of functions $\{Y_2^m, Y_2^{m*}\}$ yields a basis of a ten-dimensional (10D) space, hosting a real representation of the symmetry group $\mathbf{O}_3 \otimes \mathbf{U}_1 \times \langle \mathcal{T} \rangle$. A general element, γ , of the group will be denoted by a triple

$$\gamma = (\rho, e^{i\phi}, \epsilon),$$

where, $\rho \in \mathbf{O}_3$, $0 \leq \phi < 2\pi$ and $\epsilon = -1$, or $+1$ according as a time reflection is involved in the transformation, or not. In the following, we shall also use the shortened notations ρ for $(\rho, 1, 1)$, $U_1(\phi)$ for $(\mathbf{1}_3, U_1(\phi), 1)$ and $\mathcal{T}$ for $(\mathbf{1}_3, 1, -1)$.

The action of G can be transferred to a real irreducible action on the 10D vector formed by the coefficients $\{D_2, \dots, D_{-2}, D_2^*, \dots, D_2^*\}$. The representation of G thus obtained can be realized in the 10D real vector space generated by a couple of two independent, real, second rank, symmetric, traceless tensors, $X_{ij}^{(1)}$ and $X_{ij}^{(2)}$, $i, j = 1, 2, 3$, which can be considered as the real and imaginary parts of a complex 3×3 matrix ψ , whose elements will be written in terms of five complex coordinates, z_j :

$$z_j = x_j + i x_{5+j}, \quad j = 1, \dots, 5; \quad x_i \in \mathbf{R}, \quad (5)$$

$$\psi = \frac{1}{\sqrt{2}} \begin{pmatrix} z_2 + \frac{z_5}{\sqrt{3}} & z_1 & z_4 \\ z_1 & -z_2 + \frac{z_5}{\sqrt{3}} & z_3 \\ z_4 & z_3 & -\frac{2z_5}{\sqrt{3}} \end{pmatrix}. \quad (6)$$

The coordinates D_α are connected to the z_j by the following relations:

$$D_2 = -\frac{i z_1 + z_2}{\sqrt{2}}, \quad D_1 = \frac{i z_3 + z_4}{\sqrt{2}}, \quad D_0 = z_5, \quad D_{-1} = \frac{i z_3 - z_4}{\sqrt{2}}, \quad D_{-2} = \frac{i z_1 - z_2}{\sqrt{2}}. \quad (7)$$

The matrix ψ transforms in the following way under a general transformation, $\gamma = (\rho, \phi, \epsilon) \in G$:

$$\gamma \cdot \psi = e^{i\phi} \rho \psi' \rho^T, \quad (8)$$

where $\psi' = \psi$ or ψ^* , according as $\epsilon = +1$ or -1 and the apex T denotes transposition. As a consequence, the group G acts as a group of linear, real, *orthogonal* transformations on the vector order parameter $x = (x_1, \dots, x_{10}) \in \mathbf{R}^{10}$.

The kernel of the representation of G just defined is the group generated by the space reflection, so, it will not be restrictive to assume that the symmetry group is

²Here and in the following we denote by $\langle g_1, \dots, g_n \rangle$ the group generated by the set $\{g_1, \dots, g_n\}$.

$$G = \mathbf{SO}_3 \otimes \mathbf{U}_1 \times \langle T \rangle$$

and, when referring to G , in the following, we shall always mean this linear group acting in the vector space $\mathbf{R}^{10}$.

The group G has a trivial principal isotropy subgroup (the isotropy subgroup of generic points of $\mathbf{R}^{10}$), thus the principal G -orbits have the same dimensions, four, as G and its *orbit space*, $\mathbf{R}^{10}/G$, has dimensions $q_0 = 10 - 4 = 6$.

The Mölien function of G , $M_G(\eta)$, can be calculated in the form of an invariant Haar integral over G (see, for instance, [18,19]):

$$M_G(\eta) = \int_G \frac{d\mu(g)}{\det(\mathbf{1} - \eta g)}, \quad |\eta| < 1,$$

where $\mu(g)$ is a normalized invariant measure on the group G , the integration is over the whole group G and $g \in G$. In terms of the Euler angles ϕ_1, θ, ϕ_2 and of the gauge angle α , the integral can be written in the form of a sum over the two connected components of G

$$M_G(\eta) = \frac{1}{16\pi^2} \int_0^{2\pi} d\phi_1 \int_0^{2\pi} d\phi_2 \int_0^\pi \sin \theta d\theta \times \left\{ \int_0^{2\pi} \frac{d\alpha}{2\pi} \prod_{k=-2}^2 \left[(1 - \eta e^{i(k\chi+\alpha)}) (1 - \eta e^{i(k\chi-\alpha)}) \right]^{-1} + \prod_{k=-2}^2 \left[(1 - \eta e^{ik\chi}) (1 + \eta e^{ik\chi}) \right]^{-1} \right\}, \quad (9)$$

where $\{e^{ik\chi}\}_{-2 \leq k \leq 2}$ are the distinct eigenvalues of the 10D rotation matrix and $\cos \chi$ is a function of the Euler angles³:

$$\cos \chi = \frac{1}{2} [\cos \theta + \cos(\phi_1 + \phi_2) + \cos \theta \cos(\phi_1 + \phi_2) - 1]. \quad (10)$$

An explicit calculation of the integrals (see Appendix A) leads to

$$M_G(\eta) = \frac{1 + \eta^8 + \eta^{10} + \eta^{12} + \eta^{20}}{(1 - \eta^2)(1 - \eta^4)^2(1 - \eta^6)^2(1 - \eta^8)}, \quad (11)$$

a relation that can also be written in the following equivalent suggestive form:

$$M_G(\eta) = \frac{1 - \eta^{16} - \eta^{18} - \eta^{20} - \eta^{22} - \eta^{24} + \eta^{26} + \eta^{28} + \eta^{30} + \eta^{32} + \eta^{34} - \eta^{50}}{(1 - \eta^2)(1 - \eta^4)^2(1 - \eta^6)^2(1 - \eta^8)^2(1 - \eta^{10})(1 - \eta^{12})}. \quad (12)$$

Equations (11,12) yield the following indications, whose validity has been checked through direct calculations⁴:

1. A minimal integrity basis (IB) for the linear group G contains nine elements, $\{p_1, \dots, p_9\}$ with degrees $(d_1, \dots, d_9) = (2, 4, 4, 4, 6, 6, 8, 8, 10, 12)$.
2. The invariants p_i are connected by five independent syzygies of degrees 16, 18, 20, 22 and 24. The group is, therefore, *non coregular*.
3. It is possible to find a *non minimal* homogeneous IB, $\{\xi_1, \dots, \xi_6, \theta_1, \dots, \theta_4\}$, with degrees $(2, 4, 4, 6, 6, 8, 8, 10, 12, 20)$, such that any invariant polynomial, F , can be written in the form [20]

$$F = Q_0(\xi) + \sum_{i=1}^4 Q_i(\xi) \theta_i, \quad (13)$$

where the Q 's, are polynomials in the *algebraically independent* polynomial invariants $(\xi_1, \dots, \xi_6)$.

³It has to be noted that, in principle, the denominators in (9), are polynomials in $\sin \chi$ and $\cos \chi$; but, by they definition, they are also even functions of χ , so, they can be expressed as polynomials only in $\cos \chi$.

⁴To justify the statements listed in the following items 1)–3), starting from (11,12), one has to check that any G -invariant homogeneous polynomial of degree 20 in $x \in \mathbf{R}^{10}$ can be written as a polynomial in $p_1(x), \dots, p_9(x)$.

A possible choice for the IB's $\{p\}$ and $\{\xi, \theta\}$ is the following:

$$\begin{aligned}
p_1 &= \text{Tr}(\psi\psi^*) = \sum_{i=1}^{10} x_i^2, & p_6 &= \Re \left[\text{Tr}(\psi^2) \text{Tr}(\psi^2\psi^*) \text{Tr}(\psi^{*3}) \right], \\
p_2 &= \text{Tr} \left[(\psi\psi^*)^2 \right], & p_7 &= \Re \left[\text{Tr}(\psi^{*2}) (\text{Tr}(\psi^2\psi^*))^2 \right], \\
p_3 &= |\text{Tr}(\psi^2)|^2, & p_8 &= \Re \left[\text{Tr}(\psi^2\psi^*) \text{Tr}(\psi^3) (\text{Tr}(\psi^{*2}))^2 \right], \\
p_4 &= |\text{Tr}(\psi^3)|^2, & p_9 &= \Re \left[(\text{Tr}(\psi^2))^3 (\text{Tr}(\psi^{*3}))^2 \right], \\
p_5 &= |\text{Tr}(\psi^2\psi^*)|^2,
\end{aligned} \tag{14}$$

and

$$\begin{aligned}
\xi_i &= p_i, \quad i = 1, \dots, 5, & \xi_6 &= p_6 + p_7, \\
\theta_1 &= p_6 - p_7, & \theta_2 &= p_8, & \theta_3 &= p_9, & \theta_4 &= (p_6 - p_7)p_9.
\end{aligned} \tag{15}$$

Thus, the most general non-equilibrium polynomial Landau potential, $\widehat{\Phi}(\alpha; p)$, can be written in the form

$$\widehat{\Phi} = Q_0 + Q_1 p_7 + Q_2 p_8 + Q_3 p_9 + Q_4 p_7 p_9. \tag{16}$$

Using the above definitions, the explicit form of the syzygies and of the $\widehat{P}$ -matrix elements can be obtained through the following simple procedure.

After defining the *weight* of an element, p_i , of the IB to be equal to the degree, d_i , of the homogeneous polynomial $p_i(x)$, and the degree of $\prod_i p_i^{n_i}$ as $\sum_i n_i d_i$, one writes down the most general homogeneous polynomial, $\widehat{\Theta}^{(d)}(\xi, \theta)$, of the relevant weight, d , in the weighted variables ξ_i , $i = 1, \dots, 6$ and θ_i , $i = 1, \dots, 4$. With the definitions $\theta_0 = 1$, $d_0 = 0$, the polynomial $\widehat{\Theta}^{(d)}(\xi, \theta)$ can be written in the compact form

$$\widehat{\Theta}^{(d)}(\xi, \theta) = \sum_{j=0}^3 \delta_{N_j, d} A_{n_1, \dots, n_6}^{(j, d)} \theta_j \prod_{k=1}^6 \xi^{n_k},$$

where, $N_j = \sum_{i=1}^6 n_i d_i - d_j$, $\delta_{N_j, d}$ is a Kronecker symbol and the A 's are real coefficient to be determined through the following conditions i) and ii):

- i) in the case of the syzygies $\widehat{s}_d(\xi, \theta)$, $d = 16, 18, 20, 22, 24$: The expression $\widehat{s}_d(\xi(x), \theta(x))$ has to vanish identically in x ;
- ii) in the case of the matrix elements $\widehat{P}_{a,b}(\xi, \theta)$, $a, b = 1, \dots, 9$, $d = d_a + d_b - 2$: The expression $\widehat{P}_{ab}(\xi(x), \theta(x)) = \sum_{i=1}^{10} \partial_i p_a(x) \partial_i p_b(x)$ has to be an identity in x .

The explicit expressions of the syzygies are reported in Appendix B and the elements of the matrix $\widehat{P}(p)$ in Appendix C. In the following we shall always use the minimal IB $\{p\}$.

On the algebraic surface, Z , defined by the syzygies in the 9D space of the p 's, the 9×9 symmetric matrix $\widehat{P}(p)$ has rank 6, as it has to be, since the orbit space is six dimensional.

IV. THE ORBIT SPACE OF THE LINEAR GROUP G

As recalled in the Introduction, the orbit space of G can be identified with the range of the orbit map $x \mapsto p(x)$, which is the same as the set of values of $p \in Z$, that render the matrix $\widehat{P}(p)$ positive semi-definite.

The analytic conditions determining the geometrical strata, $\widehat{W}^{(d, \alpha, r)}$, are not difficult to specify, in principle, but, in the case of the higher dimensional strata, their explicit expressions (as sets of equalities and inequalities in the p 's) become difficult to obtain, too large to be written down extensively and uneasy to handle, if they are derived only from rank and positivity conditions of the matrix $\widehat{P}(p)$. An elegant way to bypass this difficulty is the following.

One starts with the relatively easy determination of the *geometrical* strata of dimensions ≤ 2 (i.e., the connected components of the sets $\widehat{W}^{(d)}$, $d = 1, 2$, defined in (4)), starting from the positivity and rank properties of the matrix $\widehat{P}(p)$ and from the syzygies. The subsequent step consists in the identification of the orbit-type of each geometrical stratum. This allows to identify the connected components, $\widehat{W}^{(d, \alpha, r)}$, $r = 1, \dots$, with the same orbit-type, $[G^{(d, \alpha)}]$,

and belonging, therefore, to the same *symmetry* stratum⁵, $\widehat{S}^{(d,\alpha)}$. All the groups $G^{(2,\alpha)}$ turn out to be finite groups of low order (≤ 8).

The determination of the higher dimensional strata bordered by a given 2D stratum, $\widehat{S}^{(2,\alpha)}$, can be obtained starting from the identification of their orbit-types, which are necessarily contained in the orbit-type of $\widehat{S}^{(2,\alpha)}$. The task is easily accomplished by selecting, out of the set of maximal subgroups of each $G^{(2,\alpha)}$, those which are isotropy subgroups of G . To this end, the following criterion can be used. Let H be one of the maximal subgroups of a $G^{(d,\alpha)}$ and denote by V_H the linear subspace formed by all the points $x \in \mathbf{R}^{10}$, which are stable under H . Then, H is an isotropy subgroup of G , if and only if the following condition is satisfied:

$$\text{Max}_{x \in V_H} \text{rank}(P(x)) < d.$$

If this condition is satisfied, when x spans V_H , the point $p(x) \in \mathbf{R}^9$ spans the topological closure of a stratum $\widehat{S}^{(d',\alpha')}$ and

$$\dim(\widehat{S}^{(d',\alpha')}) = \text{Max}_{x \in V_H} \text{rank}(P(x)).$$

To get the full set of higher dimensional strata, the last part of the procedure just described has to be repeated for each newly determined stratum.

The derivation of the equalities and inequalities determining each of the higher dimensional strata requires a more sophisticated method [21], which we shall explain and exploit in the following section.

Before resuming the results obtained with the realization of the program just presented, let us note that, owing to the linearity of G , its isotropy subgroups coincide at the points $x \in \mathbf{R}^{10}$ and κx , $0 \neq \kappa \in \mathbf{R}$. Thus, taking also into account the homogeneity of the polynomials $p_i(x)$, the points $(p_1, \dots, p_9)$ and $(1, p_2/p_1^{d_2/2}, \dots, p_9/p_1^{d_9/2})$ lie on the same stratum and the equalities and inequalities defining the strata can be written as homogeneous relations in the weighted variables p_i . It will be useful, therefore, to introduce the definitions

$$\tilde{p}_i = p_i/p_1^{d_i/2}, \quad \tilde{p} = (\tilde{p}_2, \dots, \tilde{p}_9). \quad (17)$$

The relations determining the 1D and 2D strata can be easily obtained by selecting those solutions, $\bar{p}$, of the condition $\text{rank}(\widehat{P}(p)) = 1$ and, respectively, $\text{rank}(\widehat{P}(p)) = 2$, at which the matrix $\widehat{P}(\bar{p})$ is positive semi-definite. In this way, one obtains five connected components, $\widehat{W}^{(1,A)}$, $A = 1, \dots, 5$, of $\widehat{W}^{(1)}$ and seven connected components, $\widehat{W}^{(2,A)}$, $A = 1_+, 1_-, 2, 3_+, 3_-, 4, 5$, of $\widehat{W}^{(2)}$. The parametric equations defining this semi-algebraic sets are listed in Tables I and II.

In order to identify the distinct 1D and 2D symmetry strata, for each connected component $\widehat{W}^{(d,A)}$, $d = 1, 2$, we have picked up a point $p^{(d,A)}$ and, for each $p^{(d,A)}$, a "simple" solution, $x^{(k,A)}$, of the equation $\tilde{p}(x) = \tilde{p}^{(d,A)}$. Then, at each point $x^{(d,A)}$ we have determined, and compared for conjugation, the isotropy subgroup of G , $G^{(d,A)}$. These subgroups turn out to be non conjugate in G , but for $G^{(2,1+)}$, which is conjugate to $G^{(2,1-)}$, and $G^{(2,3+)}$, which is conjugate to $G^{(2,3-)}$. Thus, there are five distinct 1D and five 2D symmetry strata, $\widehat{S}^{(d,\alpha)}$, with orbit-types $[G^{(d,\alpha)}]$, $d = 1, 2$, $\alpha = 1, \dots, 5$.

After choosing an element, H , in each class $[G^{(d,\alpha)}]$, it is easy to determine the subspace $V_H \subset \mathbf{R}^{10}$, formed by the vectors which are H -invariant.

The results we have obtained are listed in Tables III and IV and the bordering relations among the strata are illustrated in Figure 1. The definitions of the rotation matrices appearing in the tables and in the following text are recalled in Appendix D. We have used standard notations for the corresponding geometrical transformations (see, for instance, [22]).

⁵There is a unique zero dimensional stratum, corresponding to the point $p = p(0) = 0$, whose symmetry is $[G]$. It will be ignored in the following.

V. RELATIONS DEFINING THE STRATA WITH DIMENSIONS ≥ 3

In order to derive the relations defining the higher dimensional strata we shall exploit the following results [21], that we shall briefly recall, without proof. Let H be an isotropy subgroup of G , $S_{[H]}$ and $\hat{S}_{[H]}$ the associated strata in $\mathbf{R}^{10}$ and, respectively, $\mathbf{R}^{10}/G$ and let $\text{Stab}(H, G)$ be the stabilizer of H in G :

$$\text{Stab}(H, G) = \{g \in G \mid g H g^{-1} = H\}.$$

We have denoted by V_H the linear subspace formed by the H -invariant vectors of $\mathbf{R}^{10}$:

$$V_H = \{x \in \mathbf{R}^{10} \mid h x = x, \forall h \in H\}, \quad (18)$$

and we shall denote by $V_H^{(H)}$ the subset of V_H formed by the points at which the isotropy subgroup of G is H :

$$V_H^{(H)} = \{x \in \mathbf{R}^{10} \mid G_x = H\}. \quad (19)$$

The set $V_H^{(H)}$ is the intersection of V_H with the stratum of orbit-type $[H]$.

The group $\text{Stab}(H, G)$ acts linearly on V_H ; let us call $\tilde{H}$ the linear group defined by this action. The $\tilde{H}$ -orbit through $x \in V_H$ is the intersection of the G -orbit through x with V_H . There is, therefore, a one-to-one correspondence between the G -orbits lying in the closure of $\hat{S}_{[H]}$ and the $\tilde{H}$ orbits. This correspondence can be restricted to a one-to-one correspondence between the interior points of the closure of $\hat{S}_{[H]}$ and the points of the principal stratum of $V_H/\tilde{H}$. If $\hat{S}_{[H]}$ is connected, the one-to-one correspondence is between the principal stratum of $V_H/\tilde{H}$ and $\hat{S}_{[H]}$, otherwise the inverse image of $\hat{S}_{[H]}$ in the correspondence, reduces to the points of the principal stratum of $V_H/\tilde{H}$, corresponding to $\tilde{H}$ -orbits through points of $V_H^{(H)}$. Thus an IB, $\{\lambda\}$, of $\tilde{H}$ can be used to parameterize the points of $\hat{S}_{[H]}$:

$$p = p(\lambda).$$

The range of the parameters λ will be determined by the positivity conditions of the $\hat{P}$ -matrix associated to the IB $\{\lambda\}$, $\hat{P}^{(H)}(\lambda)$, and convenient additional conditions obtained from positivity and rank conditions of $\hat{P}(p)$, if $\hat{S}_{[H]}$ is not connected. To this end it will be worth noting that, for $p \in \hat{S}_{[H]}$, the matrix $\hat{P}(p)$ can be written in the following form:

$$\hat{P}(p(\lambda)) = J^T(\lambda) \hat{P}^{(H)}(\lambda) J(\lambda), \quad (20)$$

where, $J(\lambda)$ is the Jacobian matrix

$$J_{ai}(\lambda) = \frac{\partial p_a(\lambda)}{\partial \lambda_i}. \quad (21)$$

Owing to a well known theorem in matrix theory, from (20) one obtains the following upper limit:

$$\text{rank}(\hat{P}(p(\lambda))) \leq \text{rank}(J(\lambda)). \quad (22)$$

Below, we shall use this approach to derive rational parametric equations for all the strata of $\hat{S}$ with dimensions three and four. The results obtained in this way, starting from each of the strata of dimensions ≥ 2 , are resumed in Table IV and in the lower part of Table III.

1. Stratum $\hat{S}^{(3,1)}$

The stabilizer in G of the group $H = G^{(3,1)} = \langle C_{2z}, C_{2x}T \rangle$, is the group $\text{Stab}(H) = \langle C_{2x}, C_{2a}, U_1(\pi), T \rangle$.

An easy calculation shows that the space V_H is defined by the equations $x_i = 0$, $i = 1, 3, 4, 7, 8, 9, 10$. The elements of $\tilde{H}$ act on the x_i , $i = 2, 5, 6$ by changing the signs of one or two or all of these coordinates, so, an IB of $\tilde{H}$ can be chosen to be

$$\lambda_1 = x_5^2, \quad \lambda_2 = x_2^2, \quad \lambda_3 = x_6^2 \quad (23)$$

and the expression of the p 's in terms of the λ 's turns out to be the following:

$$\begin{aligned} p_1 &= \lambda_1 + \lambda_2 + \lambda_3, \\ p_2 &= (3\lambda_1^2 + 6\lambda_1\lambda_2 + 3\lambda_2^2 + 2\lambda_1\lambda_3 + 18\lambda_2\lambda_3 + 3\lambda_3^2)/6, \\ p_3 &= (\lambda_1 + \lambda_2 - \lambda_3)^2, \\ p_4 &= \lambda_1(\lambda_1 - 3\lambda_2 + 3\lambda_3)^2/6, \\ p_5 &= \lambda_1(\lambda_1 - 3\lambda_2 - \lambda_3)^2/6, \\ p_6 &= \lambda_1(\lambda_1 - 3\lambda_2 - \lambda_3)(\lambda_1 + \lambda_2 - \lambda_3)(\lambda_1 - 3\lambda_2 + 3\lambda_3)/6, \\ p_7 &= \lambda_1(\lambda_1 - 3\lambda_2 - \lambda_3)^2(\lambda_1 + \lambda_2 - \lambda_3)/6, \\ p_8 &= \lambda_1(\lambda_1 - 3\lambda_2 - \lambda_3)(\lambda_1 + \lambda_2 - \lambda_3)^2(\lambda_1 - 3\lambda_2 + 3\lambda_3)/6, \\ p_9 &= \lambda_1(\lambda_1 + \lambda_2 - \lambda_3)^3(\lambda_1 - 3\lambda_2 + 3\lambda_3)^2/6, \end{aligned} \quad (24)$$

with

$$\lambda_1, \lambda_2, \lambda_3 > 0. \quad (25)$$

For values of the λ 's satisfying (25), the values of the p 's defined in (24) render $\widehat{P}(p)$ everywhere positive semi-definite and of rank three. We can conclude, therefore, that they yield parametric equations for the 3D stratum $\widehat{S}^{(3,1)}$. Being the continuous image of a connected set, the stratum is *connected*.

By eliminating the parameters λ 's from (24,25), one obtains the following relations:

$$\begin{aligned} p_4 &= 9(2p_1^2 - 2p_2 - p_3)(2p_1^2 - 2p_2 + 2p_1\sqrt{p_3} + p_3)^2\tau, \\ p_5 &= (2p_1^2 - 2p_2 - p_3)(2p_1^2 - 6p_2 - 2p_1\sqrt{p_3} - p_3)^2\tau, \\ p_6 &= -3\sqrt{p_3}(2p_1^2 - 2p_2 - p_3)(2p_1^2 - 6p_2 - 2p_1\sqrt{p_3} - p_3)(2p_1^2 - 2p_2 + 2p_1\sqrt{p_3} + p_3)\tau, \\ p_7 &= -\sqrt{p_3}(2p_1^2 - 2p_2 - p_3)(2p_1^2 - 6p_2 - 2p_1\sqrt{p_3} - p_3)^2\tau, \\ p_8 &= 3p_3(2p_1^2 - 2p_2 - p_3)(2p_1^2 - 6p_2 - 2p_1\sqrt{p_3} - p_3)(2p_1^2 - 2p_2 + 2p_1\sqrt{p_3} + p_3)\tau, \\ p_9 &= -9\sqrt{p_3^3}(2p_1^2 - 2p_2 - p_3)(2p_1^2 - 2p_2 + 2p_1\sqrt{p_3} + p_3)^2\tau, \end{aligned} \quad (26)$$

$$\begin{cases} 0 < p_3 < p_1^2, \\ 2p_1^2 + p_3 < 6p_2 < 3(2p_1^2 - p_3), \end{cases} \quad (27)$$

where

$$\tau^{-1} = 64(p_1 + \sqrt{p_3})^3, \quad (28)$$

and the square root of p_3 has to be intended in an algebraic sense.

2. Stratum $\widehat{S}^{(3,2)}$

The stabilizer in G of the group $H = G^{(3,2)} = \langle C_{2x}, C_{2z} \rangle$, is a group of ∞ order: $\text{Stab}(H) = \langle C_{4x}, C_{4z}, \mathcal{T} \rangle \times \mathbf{U}_1$.

The space V_H is defined by the equations $x_i = 0$, $i = 1, 3, 4, 6, 8, 9$. The transformation properties of the coordinates x_i , $i = 2, 5, 7, 10$, under the transformations of $\widehat{H}$ can be conveniently described in the following way. Let us define:

$$\eta_1 = x_2 + ix_5 + i(x_7 + ix_{10}), \quad \eta_2 = x_2 - ix_5 + i(x_7 - ix_{10}), \quad (29)$$

then

- i) Under a gauge transformation, $\mathbf{U}_1(\phi)$: $(\eta_1, \eta_2) \rightarrow (e^{i\phi}\eta_1, e^{i\phi}\eta_2)$;
- ii) under time reversal $\mathcal{T}$: $(\eta_1, \eta_2) \rightarrow (\eta_2^*, \eta_1^*)$;
- iii) under a transformation C_{4x} : $(\eta_1, \eta_2) \rightarrow (e^{i\pi/3}\eta_2, e^{-i\pi/3}\eta_1)$;
- iv) under a transformation C_{4z} : $(\eta_1, \eta_2) \rightarrow (-\eta_2, -\eta_1)$.

The group $\tilde{H}$ is coregular and admits the following minimal IB:

$$\begin{aligned}\lambda_1 &= (|\eta_1|^2 + |\eta_2|^2)/2 = x_2^2 + x_5^2 + x_7^2 + x_{10}^2, \\ \lambda_2 &= (|\eta_1|^2 - |\eta_2|^2)^2/16 = (x_5 x_7 - x_2 x_{10})^2, \\ \lambda_3 &= \Re[(\eta_1 \eta_2^*)^3] = (x_2^2 - x_5^2 + x_7^2 - x_{10}^2) (x_2^4 - 14 x_2^2 x_5^2 + x_5^4 + 2 x_2^2 x_7^2 - \\ &\quad 2 x_5^2 x_7^2 + x_7^4 - 24 x_2 x_5 x_7 x_{10} - 2 x_2^2 x_{10}^2 + 2 x_5^2 x_{10}^2 - 14 x_7^2 x_{10}^2 + x_{10}^4).\end{aligned}\tag{30}$$

It has to be noted that the invariant $(\Im[(\eta_1 \eta_2^*)^3])^2$ can be expressed as a polynomial in $\lambda_1, \lambda_2, \lambda_3$.

The corresponding $\hat{P}$ -matrix turns out to be

$$\begin{pmatrix} 4\lambda_1 & 8\lambda_2 & 12\lambda_3 \\ 8\lambda_2 & 4\lambda_1\lambda_2 & 0 \\ 12\lambda_3 & 0 & 36\lambda_1(\lambda_1^2 - 4\lambda_2)^2 \end{pmatrix}\tag{31}$$

and the p 's can be expressed in terms of the λ 's in the following form:

$$\begin{aligned}p_1 &= \lambda_1, \\ p_2 &= (3\lambda_1^2 - 4\lambda_2)/6, \\ p_3 &= \lambda_1^2 - 4\lambda_2, \\ p_4 &= (\lambda_1^3 + 12\lambda_1\lambda_2 - \lambda_3)/12, \\ p_5 &= (\lambda_1^3 - 4\lambda_1\lambda_2 - \lambda_3)/12, \\ p_6 &= (\lambda_1^4 - 16\lambda_2^2 - \lambda_1\lambda_3)/12, \\ p_7 &= (\lambda_1^4 - 8\lambda_1^2\lambda_2 + 16\lambda_2^2 - \lambda_1\lambda_3)/12, \\ p_8 &= (\lambda_1^5 - 8\lambda_1^3\lambda_2 + 16\lambda_1\lambda_2^2 - \lambda_1^2\lambda_3 - 4\lambda_2\lambda_3)/12, \\ p_9 &= (\lambda_1^6 - 12\lambda_1^4\lambda_2 + 48\lambda_1^2\lambda_2^2 - 64\lambda_2^3 - \lambda_1^3\lambda_3 - 12\lambda_1\lambda_2\lambda_3)/12,\end{aligned}\tag{32}$$

where the parameters λ have to satisfy the conditions:

$$\begin{cases} \lambda_1 > 0, \\ 0 < 4\lambda_2 < \lambda_1^2, \\ \lambda_3^2 < (\lambda_1^2 - 4\lambda_2)^3. \end{cases}\tag{33}$$

For values of the λ 's satisfying (33), the values of the p 's defined in (32) render $\hat{P}(p)$ everywhere positive semi-definite and of rank three. We can conclude, therefore, that they yield parametric equations for the 3D stratum $\hat{S}^{(3,1)}$. Being the continuous image of a connected set, the stratum is *connected*.

By eliminating the parameters λ from (32,33) one obtains the following relations:

$$\begin{aligned}p_2 &= (2p_1^2 + p_3)/6, \\ p_5 &= (-p_1^3 + p_1p_3 + 3p_4)/3, \\ p_6 &= (-4p_1^4 + 5p_1^2p_3 - p_3^2 + 12p_1p_4)/12, \\ p_7 &= (-4p_1^4 + 3p_1^2p_3 + p_3^2 + 12p_1p_4)/12, \\ p_8 &= (-4p_1^5 + 5p_1^3p_3 - p_1p_3^2 + 12p_1^2p_4 - 6p_3p_4)/6, \\ p_9 &= (-16p_1^6 + 24p_1^4p_3 - 9p_1^2p_3^2 + p_3^3 + 48p_1^3p_4 - 36p_1p_3p_4)/12\end{aligned}\tag{34}$$

and

$$\begin{cases} p_1, p_3 > 0, \\ p_3^3 < [12p_4 - (4p_1^2 - 3p_3)p_1]^2. \end{cases}\tag{35}$$

3. Stratum $\hat{S}^{(4,1)}$

The stabilizer in G of the group $H = G^{(4,1)} = \langle C_{2x}, T \rangle$, is the group $\text{Stab}(H, G) = \mathbf{O}_2^x \times \langle U_1(\pi), T \rangle$.

The space V_H is defined by the equations $x_i = 0, i = 1, 4, 7, 8, 10$. The transformation properties of the coordinates, $x_i, i = 2, 3, 5, 6, 9$ of V_H are easily derived from (8) and can be put in the following advantageous form. Let us define

$$\eta_1 = \frac{\sqrt{3}x_2 + x_5}{2}, \quad \eta_2 = x_3 + i \frac{x_2 - \sqrt{3}x_5}{2}, \quad \eta_3 = x_6 - ix_9,\tag{36}$$

then,

- i) under a proper rotation, $O_2^x(\phi)$: $(\eta_1, \eta_2, \eta_3) \rightarrow (\eta_1, e^{-2i\phi} \eta_2, e^{i\phi} \eta_3)$;
- ii) under a gauge transformation, $U_1(\pi)$: $(\eta_1, \eta_2, \eta_3) \rightarrow (-\eta_1, -\eta_2, -\eta_3)$;
- iii) under a transformation C_{2z} : $(\eta_1, \eta_2, \eta_3) \rightarrow (\eta_1, -\eta_2^*, \eta_3^*)$;
- iv) under time reversal T : $(\eta_1, \eta_2, \eta_3) \rightarrow (\eta_1, \eta_2, -\eta_3)$.

The group $\tilde{H}$ is not coregular and admits the following minimal IB:

$$\begin{aligned}
\lambda_1 &= 3 |\eta_1|^2 = \frac{3}{4} (\sqrt{3} x_2 + x_5)^2, \\
\lambda_2 &= |\eta_2|^2 = \frac{1}{4} (x_2^2 + 4 x_3^2 - 2 \sqrt{3} x_2 x_5 + 3 x_5^2), \\
\lambda_3 &= |\eta_3|^2 = x_6^2 + x_9^2, \\
\lambda_4 &= \sqrt{3} \eta_1 \Im(\eta_2 \eta_3^2) = \frac{\sqrt{3}}{4} (\sqrt{3} x_2 + x_5) (x_2 x_6^2 - \sqrt{3} x_5 x_6^2 - 4 x_3 x_6 x_9 - \\
&\quad x_2 x_9^2 + \sqrt{3} x_5 x_9^2), \\
\lambda_5 &= [\Im(\eta_2 \eta_3^2)]^2 = \frac{1}{4} (x_2 x_6^2 - \sqrt{3} x_5 x_6^2 - 4 x_3 x_6 x_9 - x_2 x_9^2 + \sqrt{3} x_5 x_9^2)^2.
\end{aligned} \tag{37}$$

with the syzygy

$$\lambda_4^2 - \lambda_1 \lambda_5 = 0. \tag{38}$$

and the $\hat{P}$ -matrix

$$\begin{pmatrix}
12 \lambda_1 & 0 & 0 & 6 \lambda_4 & 0 \\
0 & 4 \lambda_2 & 0 & 2 \lambda_4 & 4 \lambda_5 \\
0 & 0 & 4 \lambda_3 & 4 \lambda_4 & 8 \lambda_5 \\
6 \lambda_4 & 2 \lambda_4 & 4 \lambda_4 & 4 \lambda_1 \lambda_2 \lambda_3 + \lambda_1 \lambda_3^2 + 2 \lambda_3 (4 \lambda_2 + \lambda_3) \lambda_4 & \\
0 & 4 \lambda_5 & 8 \lambda_5 & 2 \lambda_3 (4 \lambda_2 + \lambda_3) \lambda_4 & 4 \lambda_3 (4 \lambda_2 + \lambda_3) \lambda_5
\end{pmatrix} \tag{39}$$

The conditions $\hat{P}(\lambda) \geq 0$ and $\text{rank}(\hat{P}(\lambda)) = 4$, added to (38), yield the following restrictions on the acceptable range for the λ 's:

$$\begin{cases} \lambda_1, \lambda_2, \lambda_3 \geq 0, \\ 0 \leq \lambda_5 < \lambda_2 \lambda_3^2, \\ \lambda_1 + \lambda_5 > 0. \end{cases} \tag{40}$$

The p 's can be expressed in terms of the λ 's in the following form:

$$\begin{aligned}
p_1 &= (\lambda_1 + 3 \lambda_2 + 3 \lambda_3) / 3, \\
p_2 &= (\lambda_1^2 + 6 \lambda_1 \lambda_2 + 9 \lambda_2^2 + 14 \lambda_1 \lambda_3 + 18 \lambda_2 \lambda_3 + 9 \lambda_3^2 + 24 \lambda_4) / 18 \\
p_3 &= (\lambda_1 + 3 \lambda_2 - 3 \lambda_3)^2 / 9, \\
p_4 &= (4 \lambda_1^3 - 72 \lambda_1^2 \lambda_2 + 324 \lambda_1 \lambda_2^2 - 36 \lambda_1^2 \lambda_3 + 324 \lambda_1 \lambda_2 \lambda_3 + \\
&\quad 81 \lambda_1 \lambda_3^2 + 108 \lambda_1 \lambda_4 - 972 \lambda_2 \lambda_4 - 486 \lambda_3 \lambda_4 + 729 \lambda_5) / 648, \\
p_5 &= (4 \lambda_1^3 - 72 \lambda_1^2 \lambda_2 + 324 \lambda_1 \lambda_2^2 + 12 \lambda_1^2 \lambda_3 - 108 \lambda_1 \lambda_2 \lambda_3 + \\
&\quad 9 \lambda_1 \lambda_3^2 - 36 \lambda_1 \lambda_4 + 324 \lambda_2 \lambda_4 - 54 \lambda_3 \lambda_4 + 81 \lambda_5) / 648, \\
p_6 &= (\lambda_1 + 3 \lambda_2 - 3 \lambda_3) (4 \lambda_1^3 - 72 \lambda_1^2 \lambda_2 + 324 \lambda_1 \lambda_2^2 - 12 \lambda_1^2 \lambda_3 + \\
&\quad 108 \lambda_1 \lambda_2 \lambda_3 - 27 \lambda_1 \lambda_3^2 + 36 \lambda_1 \lambda_4 - 324 \lambda_2 \lambda_4 + 162 \lambda_3 \lambda_4 - 243 \lambda_5) / 1944, \\
p_7 &= (\lambda_1 + 3 \lambda_2 - 3 \lambda_3) (4 \lambda_1^3 - 72 \lambda_1^2 \lambda_2 + 324 \lambda_1 \lambda_2^2 + 12 \lambda_1^2 \lambda_3 - \\
&\quad 108 \lambda_1 \lambda_2 \lambda_3 + 9 \lambda_1 \lambda_3^2 - 36 \lambda_1 \lambda_4 + 324 \lambda_2 \lambda_4 - 54 \lambda_3 \lambda_4 + 81 \lambda_5) / 1944, \\
p_8 &= (\lambda_1 + 3 \lambda_2 - 3 \lambda_3)^2 (4 \lambda_1^3 - 72 \lambda_1^2 \lambda_2 + 324 \lambda_1 \lambda_2^2 - 12 \lambda_1^2 \lambda_3 + \\
&\quad 108 \lambda_1 \lambda_2 \lambda_3 - 27 \lambda_1 \lambda_3^2 + 36 \lambda_1 \lambda_4 - 324 \lambda_2 \lambda_4 + 162 \lambda_3 \lambda_4 - 243 \lambda_5) / 5832, \\
p_9 &= (\lambda_1 + 3 \lambda_2 - 3 \lambda_3)^3 (4 \lambda_1^3 - 72 \lambda_1^2 \lambda_2 + 324 \lambda_1 \lambda_2^2 - 36 \lambda_1^2 \lambda_3 + \\
&\quad 324 \lambda_1 \lambda_2 \lambda_3 + 81 \lambda_1 \lambda_3^2 + 108 \lambda_1 \lambda_4 - 972 \lambda_2 \lambda_4 - 486 \lambda_3 \lambda_4 + 729 \lambda_5) / 17496.
\end{aligned} \tag{41}$$

For values of the λ 's satisfying (38) and (40), the values of the p 's defined in (41) render $\hat{P}(p)$ everywhere positive semi-definite and of rank four. We can conclude, therefore, that they yield parametric equations for the 4D stratum $\hat{S}^{(4,1)}$. Being the continuous image of a connected set, the stratum is *connected*.

For $\lambda_1 \neq 0$ [$\lambda_5 \neq 0$], the syzygy can be solved with respect to λ_5 [λ_1] and the expression one obtains can be substituted into (41), so, in this region, $(\lambda_1, \dots, \lambda_4)$ [$(\lambda_2, \dots, \lambda_5)$] play the role of local coordinates for the manifold underlying the stratum. Since, owing to (40), λ_1 and λ_5 cannot vanish simultaneously, the whole stratum is covered by two rational charts of local coordinates.

The elimination of the parameters λ from (41,40) leads to cumbersome relations, which it is not worth writing down.

4. Stratum $\widehat{S}^{(4,2)}$

The stabilizer in G of the isotropy subgroup $H = G^{(4,2)} = \langle C_{2z} \rangle$, is the group $\text{Stab}(H) = \mathbf{O}_2^z \times \mathbf{U}_1$.

The space V_H is defined by the equations $x_i = 0$, $i = 3, 4, 8, 9$. The transformation properties of the coordinates, x_i , $i = 1, 2, 5, 6, 7, 10$ of V_H can be put in the following advantageous form. Let us define

$$\eta_1 = x_5 + ix_{10}, \quad \eta_2 = \frac{x_7 + ix_6 + i(x_2 + ix_1)}{\sqrt{2}}, \quad \eta_3 = \frac{x_7 + ix_6 - i(x_2 + ix_1)}{\sqrt{2}}, \quad (42)$$

then,

- i) under a proper rotation, $\mathbf{O}_2^z(\phi)$: $(\eta_1, \eta_2, \eta_3) \rightarrow (\eta_1, e^{-2i\phi}, e^{-2i\phi} \eta_3)$;
- ii) under a gauge transformation, $\mathbf{U}_1(\phi)$: $(\eta_1, \eta_2, \eta_3) \rightarrow (e^{i\phi} \eta_1, e^{-i\phi} \eta_2, e^{i\phi} \eta_3)$;
- iii) under a transformation C_{2x} : $(\eta_1, \eta_2, \eta_3) \rightarrow (\eta_1, \eta_3^*, \eta_2^*)$;
- iv) under time reversal $\mathcal{T}$: $(\eta_1, \eta_2, \eta_3) \rightarrow (\eta_1^*, -\eta_3, -\eta_2)$.

A minimal IB of $\tilde{H}$, which is coregular, can be chosen to be

$$\begin{aligned} \lambda_1 &= \sum_{j=1}^3 |\eta_j|^2 = x_1^2 + x_2^2 + x_5^2 + x_6^2 + x_7^2 + x_{10}^2, \\ \lambda_2 &= |\eta_1|^2 = x_5^2 + x_{10}^2, \\ \lambda_3 &= (|\eta_2|^2 - |\eta_3|^2)^2/4 = (x_2 x_6 - x_1 x_7)^2, \\ \lambda_4 &= 2 \Re(\eta_1^2 \eta_2 \eta_3^*) = -x_1^2 x_5^2 - x_2^2 x_5^2 + x_5^2 x_6^2 + x_5^2 x_7^2 - \\ &\quad 4 x_1 x_5 x_6 x_{10} - 4 x_2 x_5 x_7 x_{10} + x_1^2 x_{10}^2 + x_2^2 x_{10}^2 - x_6^2 x_{10}^2 - x_7^2 x_{10}^2. \end{aligned} \quad (43)$$

The $\widehat{P}$ matrix relative to the IB $\{\lambda\}$ has the following form:

$$\widehat{P}(\lambda) = 4 \begin{pmatrix} \lambda_1 & \lambda_2 & 2\lambda_3 & 2\lambda_4 \\ \lambda_2 & \lambda_2 & 0 & \lambda_4 \\ 2\lambda_3 & 0 & (\lambda_1 - \lambda_2)\lambda_3 & 0 \\ 2\lambda_4 & \lambda_4 & 0 & \lambda_2(\lambda_1^2 - \lambda_1\lambda_2 - 4\lambda_3) \end{pmatrix} \quad (44)$$

The conditions $\widehat{P}(\lambda) > 0$ and $\text{rank}(\widehat{P}(p)) = 4$ yield the following restrictions on the acceptable values of the λ 's:

$$\begin{cases} 0 < \lambda_2 < \lambda_1, \\ 0 < 4\lambda_3 < (\lambda_1 - \lambda_2)^2, \\ \lambda_4^2 < \lambda_2^2 [(\lambda_1 - \lambda_2)^2 - 4\lambda_3]. \end{cases} \quad (45)$$

The expressions of the p 's in terms of the λ 's turn out to be the following:

$$\begin{aligned}
p_1 &= \lambda_1, \\
p_2 &= (3\lambda_1^2 - 2\lambda_1\lambda_2 + 2\lambda_2^2 + 12\lambda_3 - 2\lambda_4)/6, \\
p_3 &= \lambda_1^2 - 2\lambda_1\lambda_2 + 2\lambda_2^2 - 4\lambda_3 - 2\lambda_4, \\
p_4 &= \lambda_2(9\lambda_1^2 - 18\lambda_1\lambda_2 + 10\lambda_2^2 - 36\lambda_3 + 6\lambda_4)/6, \\
p_5 &= (5\lambda_1^2\lambda_2 - 14\lambda_1\lambda_2^2 + 10\lambda_2^3 - 4\lambda_2\lambda_3 - 4\lambda_1\lambda_4 + 6\lambda_2\lambda_4)/6, \\
p_6 &= (6\lambda_1^3\lambda_2 - 17\lambda_1^2\lambda_2^2 + 14\lambda_1\lambda_2^3 - 2\lambda_2^4 - 24\lambda_1\lambda_2\lambda_3 + 20\lambda_2^2\lambda_3 - \\
&\quad 3\lambda_1^2\lambda_4 + 2\lambda_1\lambda_2\lambda_4 + 4\lambda_2^2\lambda_4 + 12\lambda_3\lambda_4 - 2\lambda_4^2)/6, \\
p_7 &= (4\lambda_1^3\lambda_2 - 11\lambda_1^2\lambda_2^2 + 6\lambda_1\lambda_2^3 + 2\lambda_2^4 - 16\lambda_1\lambda_2\lambda_3 + 28\lambda_2^2\lambda_3 - \\
&\quad 5\lambda_1^2\lambda_4 + 10\lambda_1\lambda_2\lambda_4 - 4\lambda_2^2\lambda_4 + 4\lambda_3\lambda_4 + 2\lambda_4^2)/6, \\
p_8 &= (3\lambda_1^4\lambda_2 + 2\lambda_1^3\lambda_2^2 - 36\lambda_1^2\lambda_2^3 + 52\lambda_1\lambda_2^4 - 20\lambda_2^5 - 24\lambda_1^2\lambda_2\lambda_3 - \\
&\quad 8\lambda_1\lambda_2^2\lambda_3 + 80\lambda_2^3\lambda_3 + 48\lambda_2\lambda_3^2 - 6\lambda_1^3\lambda_4 + 16\lambda_1^2\lambda_2\lambda_4 - 16\lambda_1\lambda_2^2\lambda_4 + \\
&\quad 8\lambda_2^3\lambda_4 + 24\lambda_1\lambda_3\lambda_4 - 16\lambda_2\lambda_3\lambda_4 - 4\lambda_1\lambda_4^2 + 12\lambda_2\lambda_4^2)/6, \\
p_9 &= (33\lambda_1^4\lambda_2^2 - 132\lambda_1^3\lambda_2^3 + 168\lambda_1^2\lambda_2^4 - 72\lambda_1\lambda_2^5 + 4\lambda_2^6 - 264\lambda_1^2\lambda_2^2\lambda_3 + \\
&\quad 528\lambda_1\lambda_2^3\lambda_3 - 144\lambda_2^4\lambda_3 + 528\lambda_2^2\lambda_3^2 - 9\lambda_1^4\lambda_4 + 36\lambda_1^3\lambda_2\lambda_4 - 60\lambda_1^2\lambda_2^2\lambda_4 + \\
&\quad 48\lambda_1\lambda_2^3\lambda_4 - 12\lambda_2^4\lambda_4 + 72\lambda_1^2\lambda_3\lambda_4 - 144\lambda_1\lambda_2\lambda_3\lambda_4 + 96\lambda_2^2\lambda_3\lambda_4 - \\
&\quad 144\lambda_3^2\lambda_4 - 12\lambda_1^2\lambda_4^2 + 24\lambda_1\lambda_2\lambda_4^2 + 12\lambda_2^2\lambda_4^2 + 48\lambda_3\lambda_4^2 - 4\lambda_4^3)/6.
\end{aligned} \tag{46}$$

In the range of values for the λ 's defined in (45), the values of the p 's defined in (46) render $\hat{P}(p)$ everywhere positive semi-definite and of rank four. We can conclude, therefore, that they yield parametric equations for the 4D stratum $\hat{S}^{(4,2)}$. Being the continuous image of a connected set, the stratum is *connected*.

The elimination of the parameters λ from (46,45) leads to cumbersome relations, which it is not worthwhile writing down.

5. Principal stratum $\hat{S}^{(6,1)}$

Implicit equations for the principal stratum are yielded in $\mathbf{R}^9$ by the syzygies relating the invariants p_i , $i = 1, \dots, 9$, which can be easily obtained from the syzygies (B1-B5), using (15). At the points of $\hat{S}_p = \hat{S}^{(6,1)}$ the matrix $\hat{P}(p)$ has to be positive semi-definite and its rank has to be 6.

The method used for the singular strata does not help to obtain parametric equations for $\hat{S}_p$ and to attain the goal one has to resort to less elegant and more pedantic procedures. A possibility is to use the invariants of an IB of the subgroup $\mathbf{SO}_3$ of G as parameters in terms of which to express the p 's, but in this case, great care has to be paid to the determination of the range of the parameters that render one-to-one the correspondence with the points $p \in \hat{S}_p$.

The Mölien function for the subgroup $\mathbf{SO}_3$ of G can be obtained as a byproduct of the calculation of the Mölien function for G (see Appendix A):

$$M_{\mathbf{SO}_3}(\eta) = \frac{1 + \eta^4 + \eta^8}{(1 - \eta^2 - \eta^2)^3 (1 - \eta^3)^4}, \tag{47}$$

which, after multiplying numerator and denominator by $(1 - \eta^4)$, becomes

$$M_{\mathbf{SO}_3}(\eta) = \frac{1 - \eta^{12}}{(1 - \eta^2)^3 (1 - \eta^3)^4 (1 - \eta^4)}. \tag{48}$$

Equations (47,48) indicate that a minimal IB for the subgroup $\mathbf{SO}_3$ of G is formed by eight invariants, with degrees (2, 2, 2, 3, 3, 3, 3, 4) related by only one syzygy of degree 12. A possible choice is the following:

$$\begin{aligned}
\lambda_1 &= \text{Tr}(\psi\psi^*), & \lambda_5 &= \Im[\text{Tr}(\psi^3)], \\
\lambda_2 &= \Re[\text{Tr}(\psi^2)], & \lambda_6 &= \Re[\text{Tr}(\psi^2\psi^*)], \\
\lambda_3 &= \Im[\text{Tr}(\psi^2)], & \lambda_7 &= \Im[\text{Tr}(\psi^2\psi^*)], \\
\lambda_4 &= \Re[\text{Tr}(\psi^3)], & \lambda_8 &= \text{Tr}(\psi\psi^*\psi\psi^*).
\end{aligned} \tag{49}$$

Only the first seven elements of the basis are algebraically independent. The explicit expression of the syzygy relating the λ 's is reported in Appendix B (see (B7)).

The $\hat{P}$ -matrix relative to the IB $\{\lambda_1, \dots, \lambda_8\}$ is the following and for values of the λ 's satisfying the syzygy (B7), it has rank seven:

$$\begin{aligned}
\hat{P}_{1i} &= 2 d_i^{(\lambda)} \lambda_i, & d^{(\lambda)} &= (2, 2, 2, 3, 3, 3, 3, 4), \quad i = 1, \dots, 8, \\
\hat{P}_{22} &= 4\lambda_1, & \hat{P}_{45} &= 0, \\
\hat{P}_{23} &= 0, & \hat{P}_{46} &= \frac{1}{2} (2\lambda_1 \lambda_2 + \lambda_2^2 - \lambda_3^2), \\
\hat{P}_{24} &= 6\lambda_6, & \hat{P}_{47} &= -(\lambda_1 - \lambda_2) \lambda_3, \\
\hat{P}_{25} &= 6\lambda_7, & \hat{P}_{48} &= 2 (2\lambda_1 \lambda_4 + \lambda_2 \lambda_6 - \lambda_3 \lambda_7), \\
\hat{P}_{26} &= 2 (\lambda_4 + 2\lambda_6), & \hat{P}_{55} &= \frac{3}{4} (6\lambda_1^2 - \lambda_2^2 - \lambda_3^2 - 6\lambda_8), \\
\hat{P}_{27} &= 2 (\lambda_5 - 2\lambda_7), & \hat{P}_{56} &= (\lambda_1 + \lambda_2) \lambda_3, \\
\hat{P}_{28} &= 4\lambda_1 \lambda_2, & \hat{P}_{57} &= \frac{1}{2} (2\lambda_1 \lambda_2 - \lambda_2^2 + \lambda_3^2), \\
\hat{P}_{33} &= 4\lambda_1, & \hat{P}_{58} &= 2 (2\lambda_1 \lambda_5 + \lambda_3 \lambda_6 + \lambda_2 \lambda_7), \\
\hat{P}_{34} &= -6\lambda_7, & \hat{P}_{66} &= \frac{1}{12} (2\lambda_1^2 + 8\lambda_1 \lambda_2 + 5\lambda_2^2 + 5\lambda_3^2 + 6\lambda_8), \\
\hat{P}_{35} &= 6\lambda_6, & \hat{P}_{67} &= \frac{2}{3} \lambda_1 \lambda_3, \\
\hat{P}_{36} &= 2 (\lambda_5 + 2\lambda_7), & \hat{P}_{68} &= -\frac{2}{3} (3\lambda_2 \lambda_4 + 3\lambda_3 \lambda_5 - 14\lambda_1 \lambda_6 + 2\lambda_2 \lambda_6 + 2\lambda_3 \lambda_7), \\
\hat{P}_{37} &= -2 (\lambda_4 - 2\lambda_6), & \hat{P}_{77} &= \frac{1}{12} (2\lambda_1^2 - 8\lambda_1 \lambda_2 + 5\lambda_2^2 + 5\lambda_3^2 + 6\lambda_8), \\
\hat{P}_{38} &= 4\lambda_1 \lambda_3, & \hat{P}_{78} &= \frac{2}{3} (3\lambda_3 \lambda_4 - 3\lambda_2 \lambda_5 - 2\lambda_3 \lambda_6 + 14\lambda_1 \lambda_7 + 2\lambda_2 \lambda_7), \\
\hat{P}_{44} &= \frac{3}{4} (6\lambda_1^2 - \lambda_2^2 - \lambda_3^2 - 6\lambda_8), & \hat{P}_{88} &= -\frac{8}{3} (3\lambda_1^3 - 2\lambda_4^2 - 2\lambda_5^2 + 2\lambda_6^2 + 2\lambda_7^2 - 9\lambda_1 \lambda_8).
\end{aligned} \tag{50}$$

The connection between the p 's and the λ 's can be immediately obtained from their very definitions:

$$\begin{aligned}
p_1 &= \lambda_1, \\
p_2 &= \lambda_8, \\
p_3 &= \lambda_2^2 + \lambda_3^2, \\
p_4 &= \lambda_4^2 + \lambda_5^2, \\
p_5 &= \lambda_6^2 + \lambda_7^2, \\
p_6 &= \lambda_2 \lambda_4 \lambda_6 + \lambda_3 \lambda_5 \lambda_6 - \lambda_3 \lambda_4 \lambda_7 + \lambda_2 \lambda_5 \lambda_7, \\
p_7 &= \lambda_2 \lambda_6^2 + 2 \lambda_3 \lambda_6 \lambda_7 - \lambda_2 \lambda_7^2, \\
p_8 &= \lambda_2^2 \lambda_4 \lambda_6 - \lambda_3^2 \lambda_4 \lambda_6 + 2 \lambda_2 \lambda_3 \lambda_5 \lambda_6 + 2 \lambda_2 \lambda_3 \lambda_4 \lambda_7 - \lambda_2^2 \lambda_5 \lambda_7 + \lambda_3^2 \lambda_5 \lambda_7, \\
p_9 &= \lambda_2^3 \lambda_4^2 - 3 \lambda_2 \lambda_3^2 \lambda_4^2 + 6 \lambda_2^2 \lambda_3 \lambda_4 \lambda_5 - 2 \lambda_3^3 \lambda_4 \lambda_5 - \lambda_2^3 \lambda_5^2 + 3 \lambda_2 \lambda_3^2 \lambda_5^2.
\end{aligned} \tag{51}$$

As already noted, there is not one-to-one correspondence between the points $\lambda = (\lambda_1, \dots, \lambda_8)$, lying on the surface determined by the syzygy (B7) and rendering positive semi-definite and of rank seven the $\hat{P}$ -matrix associated to the integrity basis $\{\lambda\}$, and the points p of the principal stratum of $\mathbf{R}^{10}/G$. In fact, the action of the group G in $\mathbf{R}^{10}$ induces a linear action of G on the variables λ_i . With the definitions

$$\eta_1 = \lambda_2 + i \lambda_3, \quad \eta_2 = \lambda_4 + i \lambda_5, \quad \eta_3 = \lambda_6 + i \lambda_7, \tag{52}$$

the variables λ and η have the following transformation properties:

- i) λ_1 and λ_8 are invariant under the whole group;
- ii) under gauge transformations, $U_1(\phi)$: $(\eta_1, \eta_2, \eta_3) \rightarrow (e^{2i\phi} \eta_1, e^{3i\phi} \eta_2, e^{i\phi} \eta_3)$;
- iii) under time reversal T : $(\eta_1, \eta_2, \eta_3) \rightarrow (\eta_1^*, \eta_2^*, \eta_3^*)$.

The one-to-one correspondence problem can be solved by finding a criterion enabling to select a unique point in each orbit of the action of G in the λ space. To this end, let us make preliminary remarks;

- i) A direct check shows that for $\eta_1 = 0$ and general values of η_2 and η_3 the rank of $\hat{P}(p(\lambda))$ is six.
- ii) Using (22) one easily realizes that $\text{rank}(\hat{P}(p(\lambda))) < \text{rank}(J(\lambda)) < 6$, for $\lambda_3 = \lambda_5 = \lambda_7 = 0$.

Now, let us choose any orbit, $\bar{\Omega}$, in the principal stratum of $\mathbf{R}^{10}/G$, *i.e.*, a $\bar{p} \in \hat{S}_p$, and let $\bar{\lambda}$ be such that $p(\bar{\lambda}) = \bar{p}$. At least one of the parameters $\bar{\eta}_i$, $i = 1, 2, 3$, has to be different from zero. Let us say that $\bar{\eta}_i$ is the first one, in lexicographic order⁶. Then, on $\bar{\Omega}$ there will be a point $\bar{\lambda}'$, such that $\bar{\eta}'_i$ is real and positive; this point can be attained

⁶In a lexicographic ordering, $(v_1, \dots, v_k) > (v'_1, \dots, v'_k)$ means that the first non vanishing component $v_i - v'_i$, $i = 1, \dots, k$, is positive.

from $\bar{\lambda}$ by means of a convenient gauge transformation. Since at least a $\bar{\eta}_j$, $j = 1, 2, 3$, has to be $\neq 0$, this means $\bar{i} < 3$. At this point we still have at our disposal time reversal transformations to fix the sign of the imaginary part of one of the $\bar{\eta}_i$, $i > \bar{i}$ (recall that at least one of the $\Im(\bar{\eta}_j)$ has to be $\neq 0$). So, if $\bar{j}$ is such that $\Im(\bar{\eta}_{\bar{j}})$ is the first (in lexicographic order) non vanishing imaginary part of the $\bar{\eta}$'s, we can choose $\Im(\bar{\eta}_{\bar{j}}) > 0$.

After introducing a lexicographic order in the real vector spaces generated by the vectors formed, respectively, with the real and imaginary parts of the η 's, the criterion that we have devised can be resumed in the following simple additional conditions on these vectors:

$$\lambda_3 = 0, \quad (\lambda_2, \lambda_4, \lambda_6) > 0, \quad (\lambda_3, \lambda_5, \lambda_7) > 0, \quad \text{and } \lambda_5 = 0 \text{ for } \lambda_2 = 0. \quad (53)$$

Using these conditions, the expressions (51) of the p 's can be simplified as follows:

$$\begin{aligned} p_1 &= \lambda_1, & p_6 &= \lambda_2 (\lambda_4 \lambda_6 + \lambda_5 \lambda_7), \\ p_2 &= \lambda_8, & p_7 &= \lambda_2 (\lambda_6^2 - \lambda_7^2), \\ p_3 &= \lambda_2^2, & p_8 &= \lambda_2^2 (\lambda_4 \lambda_6 - \lambda_5 \lambda_7), \\ p_4 &= \lambda_4^2 + \lambda_5^2, & p_9 &= \lambda_3 (\lambda_4^2 - \lambda_5^2), \\ p_5 &= \lambda_6^2 + \lambda_7^2, \end{aligned} \quad (54)$$

where the λ 's are required to satisfy the syzygy (B7), the conditions rendering the matrix $\hat{P}(p)$ positive semi-definite and of rank six and $(\lambda_5, \lambda_7) > 0$.

VI. AN EXAMPLE: MINIMA OF A GENERAL FOURTH DEGREE LANDAU POLYNOMIAL

As a simple example, we have also calculated the minimum of a general fourth degree polynomial free energy

$$\hat{\Phi}^{(4)}(p) = \frac{\alpha_0}{2} p_1^2 + \sum_{j=1}^3 \alpha_j p_j, \quad \alpha_i \in \mathbf{R}; \quad i = 0, \dots, 3, \quad (55)$$

in the additional assumptions that it is bounded below and has a local maximum at the origin ($\alpha_1 < 0$). In (55) the α 's are phenomenological parameters.

Recalling the definition $\tilde{p}_i = p_i/p_1^{d_i/2}$, the polynomial $\hat{\Phi}^{(4)}(p)$ can be put in the following form:

$$\hat{\Phi}^{(4)}(p) = \frac{p_1^2}{2} \Delta + \alpha_1 p_1, \quad (56)$$

where we have defined

$$\Delta = \alpha_0 + 2\alpha_2 \tilde{p}_2 + 2\alpha_3 \tilde{p}_3. \quad (57)$$

Since, owing to its definition (14), p_1 ranges over the whole non negative real numbers, in the orbit space, the polynomial $\hat{\Phi}^{(4)}(p)$ is bounded below (in the assumption $\alpha_1 < 0$) if and only if the minimum of Δ , δ , evaluated in the section $p_1 = 1$ of the orbit space, is positive. Being the minimum of the r.h.s. of (56), thought of as a function only of $p_1 \geq 0$, equal to $-\alpha_1/(2\Delta)$, the absolute minimum of $\hat{\Phi}^{(4)}(p)|_{p \in S}$ is $-\alpha_1^2/(2\delta)$. In this way, the determination of the minimum of $\hat{\Phi}^{(4)}(p)$ is reduced to the calculation of δ .

The absolute minimum of δ in each singular stratum can be easily computed using the equations of the strata. For the principal stratum the determination of the minimum of $\hat{\Phi}^{(4)}(p)|_{p_1=1}$ as a constrained minimum is difficult and it is easier to solve equation (2) and to check subsequently if the solutions lie in the principal stratum. A comparison of the values of the minima in the different strata, in order to determine the absolute minimum, leads to the results resumed in Table V and illustrated in Figure 2, where the denomination of the strata introduced in the text has been used. Owing to the low degree of the polynomial defining $\hat{\Phi}^{(4)}(p)$ in (55), and the consequent low number of free parameters α , the absolute minimum exhibits strong degeneracy, particularly for special values of the α 's. If these special values are excluded, spontaneous breaking of the symmetry can generate only five distinct phases out of fifteen permitted in the G -symmetry; some of them are unstable. For no non trivial values of (α_2, α_3) does the absolute minimum lie on the stratum $S^{(2,2)}$.

For general values of the α 's, our results are in agreement with Mermin's ones [11]. Let us add a few words about the perturbative stability of the three degenerate phases in the region $\mathcal{R}_2$ (see Table V). For $(\alpha_0, \alpha_2, \alpha_3) \in \mathcal{R}_2$, the addition to the free energy, $\Phi^{(4)}$, of a "small" perturbation, consisting in an invariant polynomial of degree six:

$$\Theta^{(6)} = \alpha_4 p_4 + \alpha_5 p_5, \quad (58)$$

splits the three degenerate minima determined by the 4-th degree term⁷. This is easy to check, at least in the additional assumption that the perturbation leaves the absolute minimum in one of the strata corresponding to the degenerate phases. In fact, at the first perturbative order, one obtains from Tables I and II the following shifts, $\Theta_{(d,r)}^{(6)}$, in the values of the 6-th order free energy at the points where $\Phi^{(4)}(p)$ takes on its degenerate absolute minimum under consideration:

$$\Theta_{(1,2)}^{(6)} = \frac{\alpha_1^3}{\delta^3} \frac{\alpha_4 + \alpha_5}{6}, \quad \Theta_{(1,3)}^{(6)} = 0, \quad \Theta_{(2,4)}^{(6)} = \frac{\alpha_1^3}{\delta^3} (\alpha_4 + \alpha_5) \xi, \quad (59)$$

where $0 < \xi < 1/6$.

Since $-\alpha_1/\delta > 0$, the absolute minimum will be perturbatively stable on $\widehat{S}^{(1,2)}$ or, respectively, $\widehat{S}^{(1,3)}$, according as $\alpha_4 + \alpha_5$ is negative or positive.

As stressed in the Introduction, the difficulties mentioned above can be overcome if one puts less restrictive upper limits to the degree of the polynomial describing the free energy. It is trivial, for instance, to realize that the following class of bounded below polynomial functions [16] have a vanishing maximum at the origin of $\mathbf{R}^{10}$ and display an absolute minimum at the arbitrarily chosen point $\bar{p} \in \widehat{S}$:

$$\sum_{i=1}^9 \alpha_i \left[(p_i - \bar{p}_i)^{2n_i} - \bar{p}_i^{2n_i} \right], \quad (60)$$

where the α 's are positive constants and the n 's are positive integers.

The physical implications of our results and the derivation of a more realistic form of the free energy will be discussed in forthcoming papers. We shall limit ourselves, here, to note that, the D-wave content in terms, *e.g.*, of d- or extended s-waves (mentioned in the introduction) can be directly read off the fourth column of Table III, for each of the possible symmetry phases of the system under consideration.

VII. CONCLUSIONS

Through a detailed and rigorous analysis, we have determined and characterized the possible ground states of D-wave condensates in isotropic space, which are relevant both in high T_c superconductivity and in ^3He phase transitions. The problem has been formulated and solved in the framework of geometric invariant theory, that, as noted long ago [14], is the natural mathematical setting in which to deal with minimization of invariant potentials, which are plagued by degeneracy problems. This enabled us to complete and/or correct results previously obtained by other authors, using different methods.

Until now, the kind of approach we have used had only been applied to relatively simple models, with coregular symmetry groups, in condensed matter [23] and in elementary particle physics [24]. In this paper, we have fully characterized the geometry of the orbit space of the linear group $\mathbf{SO}_3 \otimes \mathbf{U}_1 \times \langle T \rangle$, acting in the space $\mathbf{R}^{10}$, a highly non trivial group, with a non coregular minimal integrity basis, formed by nine elements with degrees up to 12. For this group we have determined the explicit form of the elements of an integrity basis and of the syzygies relating these elements, the equations and inequalities determining the orbit space and its stratification and the corresponding isotropy subgroups chains. The minima of a general fourth degree polynomial invariant free energy have been recalculated and the sixth degree polynomial perturbative corrections to its absolute minimum have been determined. Some of our conclusions have been compared with previous results obtained by other authors.

To overcome some computational difficulties in the full characterization of the stratification of the orbit space, difficulties originating from the high dimensionality of the problem, we had to devise a sophisticated procedure, allowing to obtain *rational* parametric equations also for the higher dimensional singular strata. The procedure is absolutely general and is reminiscent, but in a different context and in higher dimensions, of classical techniques for parameterizing plane algebraic curves [25].

⁷The inclusion of terms of 6-th degree, depending only on p_1, p_2 and p_3 would be useless for splitting the degenerate minima, so these terms will be neglected, with no loss of generality.

ACKNOWLEDGMENTS

This paper is partially supported by RFBR, INFN and MURST

APPENDIX A: CALCULATION OF MÖLIEN FUNCTIONS

In this Appendix, we shall explicitly calculate the two integrals, I_1 and I_2 , appearing in the expression of $M_G(\eta)$ (see (9)):

$$\begin{aligned} I_1 &= \frac{1}{32\pi^3} \int_0^{2\pi} d\phi_1 \int_0^{2\pi} d\phi_2 \int_0^\pi \sin \theta d\theta \int_0^{2\pi} \frac{d\alpha}{f_1(\cos \chi, \alpha)}, \\ I_2 &= \frac{1}{16\pi^2} \int_0^{2\pi} d\phi_1 \int_0^{2\pi} d\phi_2 \int_0^\pi \frac{\sin \theta d\theta}{f_2(\cos \chi)}, \end{aligned} \tag{A1}$$

where

$$\begin{aligned} f_1(\cos \chi, \alpha) &= \prod_{k=-2}^2 \left[\left(1 - \eta e^{i(k\chi + \alpha)} \right) \left(1 - \eta e^{i(k\chi - \alpha)} \right) \right] \\ f_2(\cos \chi) &= \prod_{k=-2}^2 \left[\left(1 - \eta e^{i k \chi} \right) \left(1 + \eta e^{i k \chi} \right) \right] \end{aligned} \tag{A2}$$

and $\cos \chi$ is defined in (10).

Using as new integration variables ϕ_2 , $u = \cos(\phi_1 + \phi_2)$ and $v = \cos \chi$, the integral over ϕ_2 can be immediately performed and one remains with

$$\begin{aligned} I_1 &= \frac{1}{2\pi^2} \int_0^{2\pi} d\alpha \int_{-1}^1 \frac{dv}{f_1(v, \alpha)} \int_v^1 \frac{du}{(1+u)\sqrt{1-u^2}} \\ &= \frac{1}{2\pi^2} \int_0^{2\pi} d\alpha \int_{-1}^1 \frac{dv}{f_1(v, \alpha)} \frac{\sqrt{1-v^2}}{1+v} \\ &= \frac{1}{2\pi^2} \int_0^{2\pi} d\alpha \int_0^{2\pi} d\chi \frac{\sin^2(\chi/2)}{f_1(\cos \chi, \alpha)} \end{aligned} \tag{A3}$$

and, analogously,

$$I_2 = \frac{1}{\pi} \int_0^{2\pi} d\chi \frac{\sin^2(\chi/2)}{f_2(\cos \chi)}. \tag{A4}$$

After defining

$$\zeta_1 = e^{i\chi}, \quad \zeta_2 = e^{i\alpha}, \tag{A5}$$

the remaining integrals can be conveniently thought of as integrals on unit circles of complex planes of the variables ζ_1 and ζ_2

$$\begin{aligned} I_1 &= -\frac{1}{8\pi^2 \eta^4} \oint \frac{d\zeta_2}{(\zeta_2 - \eta)(\zeta_2 \eta - 1)} \oint d\zeta_1 F_1(\zeta_1, \zeta_2, \eta) \\ I_2 &= \frac{1}{4\pi i \eta^4 (\eta^2 - 1)} \oint d\zeta_1 F_2(\zeta_1, \eta), \end{aligned} \tag{A6}$$

where

$$\begin{aligned}
F_1(\zeta_1, \zeta_2, \eta) &= \frac{\zeta_1^4 (\zeta_1 - 1)^2}{\left(\zeta_1^2 - \frac{\eta}{\zeta_2}\right) (\zeta_1^2 - \zeta_2 \eta) \left(\zeta_1 - \frac{\eta}{\zeta_2}\right) (\zeta_1 - \zeta_2 \eta) \left(\zeta_1 - \frac{\zeta_2}{\eta}\right) \left(\zeta_1 - \frac{1}{\zeta_2 \eta}\right) \left(\zeta_1^2 - \frac{\zeta_2}{\eta}\right) \left(\zeta_1^2 - \frac{1}{\zeta_2 \eta}\right)} \\
F_2(\zeta_1, \eta) &= \frac{\zeta_1^4 (\zeta_1 - 1)^2}{(\zeta_1^2 - \eta^2) (\zeta_1^4 - \eta^2) \left(\zeta_1^2 - \frac{1}{\eta^2}\right) \left(\zeta_1^4 - \frac{1}{\eta^2}\right)}.
\end{aligned} \tag{A7}$$

Using the Cauchy theorem and recalling that $|\eta| < 1$, the calculation reduces to a computation of residues at the internal poles to the unit circles. So, one ends up with the following expressions for I_1 and I_2 :

$$\begin{aligned}
I_1 &= \frac{1 + 3\eta^8 + 2\eta^{10} + 3\eta^{12} + \eta^{20}}{2(1 - \eta^2)^3 (1 - \eta^4)^2 (1 - \eta^8) (1 - \eta + \eta^2)^2 (1 + \eta + \eta^2)^2}, \\
I_2 &= \frac{1 - \eta^2 + \eta^4}{2(1 - \eta^2)^3 (1 - \eta^4) (1 - \eta + \eta^2) (1 + \eta + \eta^2)}.
\end{aligned} \tag{A8}$$

The sum of I_1 and I_2 gives to the expression of $M_G(\eta)$ reported in (11).

As a byproduct of the calculation, one finds also the following expression for the Mölien function of the subgroup $\mathbf{SO}_3$ of G :

$$\begin{aligned}
M_{\mathbf{SO}_3}(\eta) &= \frac{1}{8\pi^2} \int_0^{2\pi} d\phi_1 \int_0^{2\pi} d\phi_2 \int_0^\pi d\theta \sin \theta f_1(\cos \chi, 0) \\
&= -\frac{1}{4\pi i \eta^4 (1 - \eta^2)} \oint d\zeta_1 F_1(\zeta_1, 1, \eta) \\
&= \frac{1 + \eta^4 + \eta^8}{(1 - \eta^2)^3 (1 - \eta^3)^4}.
\end{aligned} \tag{A9}$$

APPENDIX B: SYZYGIES

1. Explicit form of the syzygies for the group G

The ten elements of the non minimal IB $\{\xi, \theta\}$ defined in (14,15) are related by the following syzygies:

$$\begin{aligned}
\theta_1^2 &= \frac{1}{192} [36\xi_1^6 \xi_3 - 144\xi_1^4 \xi_2 \xi_3 + 180\xi_1^2 \xi_2^2 \xi_3 - 72\xi_2^3 \xi_3 - 36\xi_1^4 \xi_3^2 + 108\xi_1^2 \xi_2 \xi_3^2 - \\
&\quad 72\xi_2^2 \xi_3^2 + 9\xi_1^2 \xi_3^3 - 18\xi_2 \xi_3^3 - 80\xi_1^3 \xi_3 \xi_4 + 144\xi_1 \xi_2 \xi_3 \xi_4 + 24\xi_1 \xi_3^2 \xi_4 + 48\xi_3 \xi_4^2 + \\
&\quad 144\xi_1^3 \xi_3 \xi_5 - 144\xi_1 \xi_2 \xi_3 \xi_5 + 72\xi_1 \xi_3^2 \xi_5 - 288\xi_3 \xi_4 \xi_5 - 144\xi_3 \xi_5^2 - 96\xi_1^2 \xi_3 \xi_6 - \\
&\quad 24\xi_3^2 \xi_6 + 192\xi_6^2 + 24\theta_1 (8\xi_1^2 - 12\xi_2 - \xi_3) \xi_3 + 96\theta_2 (\xi_1 \xi_3 - 4\xi_5) - 16\theta_3 \xi_3];
\end{aligned} \tag{B1}$$

$$\theta_1 \theta_2 = -\frac{1}{2} (\xi_3 \xi_4 \xi_6 + \theta_1 \xi_3 \xi_4 - 2\theta_2 \xi_6 + 2\theta_3 \xi_5); \tag{B2}$$

$$\begin{aligned}
\theta_2^2 &= \frac{1}{768} [-36\xi_1^6 \xi_3^2 + 144\xi_1^4 \xi_2 \xi_3^2 - 180\xi_1^2 \xi_2^2 \xi_3^2 + 72\xi_2^3 \xi_3^2 + 36\xi_1^4 \xi_3^3 - \\
&\quad 108\xi_1^2 \xi_2 \xi_3^3 + 72\xi_2^2 \xi_3^3 - 9\xi_1^2 \xi_3^4 + 18\xi_2 \xi_3^4 + 80\xi_1^3 \xi_3^2 \xi_4 - 144\xi_1 \xi_2 \xi_3^2 \xi_4 - \\
&\quad 24\xi_1 \xi_3^3 \xi_4 - 48\xi_3^2 \xi_4^2 - 44\xi_1^3 \xi_3^2 \xi_5 + 144\xi_1 \xi_2 \xi_3^2 \xi_5 - 72\xi_1 \xi_3^3 \xi_5 + \\
&\quad 1056\xi_3^2 \xi_4 \xi_5 + 144\xi_3^2 \xi_5^2 + 96\xi_1^2 \xi_3^2 \xi_6 + 24\xi_3^3 \xi_6 - 384\xi_3 \xi_6^2 + \\
&\quad 24\theta_1 \xi_3 (-8\xi_1^2 \xi_3 + 12\xi_2 \xi_3 + \xi_3^2 - 16\xi_6) - 96\theta_2 \xi_3 (\xi_1 \xi_3 - 4\xi_5) + \\
&\quad 16\theta_3 (\xi_3^2 + 24\xi_6) - 384\theta_4];
\end{aligned} \tag{B3}$$

$$\begin{aligned}
\theta_2 \theta_3 = & \frac{1}{128} [-36 \xi_1^7 \xi_3^2 + 144 \xi_1^5 \xi_2 \xi_3^2 - 180 \xi_1^3 \xi_2^2 \xi_3^2 + 72 \xi_1 \xi_2^3 \xi_3^2 + 36 \xi_1^5 \xi_3^3 - \\
& 108 \xi_1^3 \xi_2 \xi_3^3 + 72 \xi_1 \xi_2^2 \xi_3^3 - 9 \xi_1^3 \xi_3^4 + 18 \xi_1 \xi_2 \xi_3^4 + 144 \xi_1^6 \xi_3 \xi_4 - 576 \xi_1^4 \xi_2 \xi_3 \xi_4 + \\
& 720 \xi_1^2 \xi_2^2 \xi_3 \xi_4 - 288 \xi_2^3 \xi_3 \xi_4 - 64 \xi_1^4 \xi_3^2 \xi_4 + 288 \xi_1^2 \xi_2 \xi_3^2 \xi_4 - 288 \xi_2^2 \xi_3^2 \xi_4 + \\
& 12 \xi_1^2 \xi_3^3 \xi_4 - 72 \xi_2 \xi_3^3 \xi_4 - 320 \xi_1^3 \xi_3 \xi_4^2 + 576 \xi_1 \xi_2 \xi_3 \xi_4^2 + 48 \xi_1 \xi_3^2 \xi_4^2 + 192 \xi_3 \xi_4^3 + \\
& 144 \xi_1^6 \xi_3 \xi_5 - 576 \xi_1^4 \xi_2 \xi_3 \xi_5 + 720 \xi_1^2 \xi_2^2 \xi_3 \xi_5 - 288 \xi_2^3 \xi_3 \xi_5 - 288 \xi_1^4 \xi_3^2 \xi_5 + \\
& 576 \xi_1^2 \xi_2 \xi_3^2 \xi_5 - 288 \xi_2^2 \xi_3^2 \xi_5 - 36 \xi_1^2 \xi_3^3 \xi_5 - 72 \xi_2 \xi_3^3 \xi_5 + 256 \xi_1^3 \xi_3 \xi_4 \xi_5 + \\
& 1440 \xi_1 \xi_3^2 \xi_4 \xi_5 - 960 \xi_3 \xi_4^2 \xi_5 + 576 \xi_1^3 \xi_3 \xi_5^2 - 576 \xi_1 \xi_2 \xi_3 \xi_5^2 + 432 \xi_1 \xi_3^2 \xi_5^2 - \\
& 4800 \xi_3 \xi_4 \xi_5^2 - 576 \xi_3 \xi_5^3 + 96 \xi_1^3 \xi_3^2 \xi_6 + 24 \xi_1 \xi_3^3 \xi_6 + 384 \xi_1^2 \xi_3 \xi_4 \xi_6 - 1152 \xi_2 \xi_3 \xi_4 \xi_6 - \\
& 192 \xi_3^2 \xi_4 \xi_6 - 384 \xi_1^2 \xi_3 \xi_5 \xi_6 - 96 \xi_3^2 \xi_5 \xi_6 - 384 \xi_1 \xi_3 \xi_6^2 + 1536 \xi_4 \xi_6^2 + 1536 \xi_5 \xi_6^2 - \\
& 24 \theta_1 (8 \xi_1^3 \xi_3^2 - 12 \xi_1 \xi_2 \xi_3^2 - \xi_1 \xi_3^3 - 32 \xi_1^2 \xi_3 \xi_5 + 48 \xi_2 \xi_3 \xi_5 + 4 \xi_3^2 \xi_5 + 16 \xi_1 \xi_3 \xi_6 + \\
& 64 \xi_4 \xi_6 - 64 \xi_5 \xi_6) + \\
& 8 \theta_2 (36 \xi_1^6 - 144 \xi_1^4 \xi_2 + 180 \xi_1^2 \xi_2^2 - 72 \xi_2^3 - 36 \xi_1^4 \xi_3 + 108 \xi_1^2 \xi_2 \xi_3 - 72 \xi_2^2 \xi_3 - \\
& 3 \xi_1^2 \xi_3^2 - 18 \xi_2 \xi_3^2 - 80 \xi_1^3 \xi_4 + 144 \xi_1 \xi_2 \xi_4 + 72 \xi_1 \xi_3 \xi_4 + 48 \xi_4^2 + 144 \xi_1^3 \xi_5 - \\
& 144 \xi_1 \xi_2 \xi_5 + 168 \xi_1 \xi_3 \xi_5 - 480 \xi_4 \xi_5 - 336 \xi_5^2 - 288 \xi_1^2 \xi_6 + 288 \xi_2 \xi_6) + \\
& 16 \theta_3 (\xi_1 \xi_3^2 - 4 \xi_3 \xi_4 + 96 \xi_1^2 \xi_5 - 144 \xi_2 \xi_5 - 16 \xi_3 \xi_5 + 24 \xi_1 \xi_6) - 384 \theta_4 \xi_1];
\end{aligned} \tag{B4}$$

$$\begin{aligned}
\theta_3^2 = & \frac{1}{256} [-144 \xi_1^8 \xi_3^2 + 144 \xi_1^6 \xi_2 \xi_3^2 + 1008 \xi_1^4 \xi_2^2 \xi_3^2 - 1872 \xi_1^2 \xi_2^3 \xi_3^2 + 864 \xi_2^4 \xi_3^2 + \\
& 108 \xi_1^6 \xi_3^3 + 144 \xi_1^4 \xi_2 \xi_3^3 - 1188 \xi_1^2 \xi_2^2 \xi_3^3 + 936 \xi_2^3 \xi_3^3 - 144 \xi_1^2 \xi_2 \xi_3^4 + 288 \xi_2^2 \xi_3^4 - \\
& 9 \xi_1^2 \xi_3^5 + 18 \xi_2 \xi_3^5 + 1728 \xi_1^7 \xi_3 \xi_4 - 6912 \xi_1^5 \xi_2 \xi_3 \xi_4 + 8640 \xi_1^3 \xi_2^2 \xi_3 \xi_4 - \\
& 3456 \xi_1 \xi_2^3 \xi_3 \xi_4 - 1408 \xi_1^5 \xi_3^2 \xi_4 + 5568 \xi_1^3 \xi_2 \xi_3^2 \xi_4 - 5184 \xi_1 \xi_2^2 \xi_3^2 \xi_4 + 416 \xi_1^3 \xi_3^3 \xi_4 - \\
& 1296 \xi_1 \xi_2 \xi_3^3 \xi_4 - 24 \xi_1 \xi_3^4 \xi_4 - 3840 \xi_1^4 \xi_3 \xi_4^2 + 6912 \xi_1^2 \xi_2 \xi_3 \xi_4^2 + 960 \xi_1^2 \xi_3^2 \xi_4^2 - \\
& 576 \xi_2 \xi_3^2 \xi_4^2 - 48 \xi_3^3 \xi_4^2 + 2304 \xi_1 \xi_3 \xi_4^3 + 1728 \xi_1^7 \xi_3 \xi_5 - 6912 \xi_1^5 \xi_2 \xi_3 \xi_5 + \\
& 8640 \xi_1^3 \xi_2^2 \xi_3 \xi_5 - 3456 \xi_1 \xi_2^3 \xi_3 \xi_5 - 2304 \xi_1^5 \xi_3^2 \xi_5 + 4032 \xi_1^3 \xi_2 \xi_3^2 \xi_5 - \\
& 1728 \xi_1 \xi_2^2 \xi_3^2 \xi_5 - 1584 \xi_1 \xi_2 \xi_3^3 \xi_5 - 72 \xi_1 \xi_3^4 \xi_5 + 3072 \xi_1^4 \xi_3 \xi_4 \xi_5 + 14976 \xi_1^2 \xi_3^2 \xi_4 \xi_5 + \\
& 3456 \xi_2 \xi_3^2 \xi_4 \xi_5 + 288 \xi_3^3 \xi_4 \xi_5 - 11520 \xi_1 \xi_3 \xi_4^2 \xi_5 + 6912 \xi_1^4 \xi_3 \xi_5^2 - 6912 \xi_1^2 \xi_2 \xi_3 \xi_5^2 + \\
& 4032 \xi_1^2 \xi_3^2 \xi_5^2 + 1728 \xi_2 \xi_3^2 \xi_5^2 + 144 \xi_3^3 \xi_5^2 - 57600 \xi_1 \xi_3 \xi_4 \xi_5^2 - 6912 \xi_1 \xi_3 \xi_5^3 + \\
& 864 \xi_1^6 \xi_3 \xi_6 - 3456 \xi_1^4 \xi_2 \xi_3 \xi_6 + 4320 \xi_1^2 \xi_2^2 \xi_3 \xi_6 - 1728 \xi_2^3 \xi_3 \xi_6 - 480 \xi_1^4 \xi_3^2 \xi_6 + \\
& 3744 \xi_1^2 \xi_2 \xi_3^2 \xi_6 - 1728 \xi_2^2 \xi_3^2 \xi_6 + 408 \xi_1^2 \xi_3^3 \xi_6 - 144 \xi_2 \xi_3^3 \xi_6 + 24 \xi_3^4 \xi_6 + \\
& 2688 \xi_1^3 \xi_3 \xi_4 \xi_6 - 10368 \xi_1 \xi_2 \xi_3 \xi_4 \xi_6 - 1344 \xi_1 \xi_3^2 \xi_4 \xi_6 - 384 \xi_3 \xi_4^2 \xi_6 - 1152 \xi_1^3 \xi_3 \xi_5 \xi_6 - \\
& 3456 \xi_1 \xi_2 \xi_3 \xi_5 \xi_6 + 576 \xi_1 \xi_3^2 \xi_5 \xi_6 - 14592 \xi_3 \xi_4 \xi_5 \xi_6 - 3456 \xi_3 \xi_5^2 \xi_6 - 5376 \xi_1^2 \xi_3 \xi_6^2 - \\
& 2304 \xi_2 \xi_3 \xi_6^2 - 768 \xi_3^2 \xi_6^2 + 18432 \xi_1 \xi_4 \xi_6^2 + 18432 \xi_1 \xi_5 \xi_6^2 + 6144 \xi_6^3 + \\
& 8 \theta_1 (36 \xi_1^6 \xi_3 - 144 \xi_1^4 \xi_2 \xi_3 + 180 \xi_1^2 \xi_2^2 \xi_3 - 72 \xi_2^3 \xi_3 - 132 \xi_1^4 \xi_3^2 - 36 \xi_1^2 \xi_2 \xi_3^2 + \\
& 360 \xi_2^2 \xi_3^2 - 3 \xi_1^2 \xi_3^3 + 54 \xi_2 \xi_3^3 + 3 \xi_3^4 - 80 \xi_1^3 \xi_3 \xi_4 + 144 \xi_1 \xi_2 \xi_3 \xi_4 - 24 \xi_1 \xi_3^2 \xi_4 + \\
& 240 \xi_3 \xi_4^2 + 1296 \xi_1^3 \xi_3 \xi_5 - 1872 \xi_1 \xi_2 \xi_3 \xi_5 - 72 \xi_1 \xi_3^2 \xi_5 - 864 \xi_3 \xi_4 \xi_5 - 144 \xi_3 \xi_5^2 - \\
& 96 \xi_1^2 \xi_3 \xi_6 - 864 \xi_2 \xi_3 \xi_6 - 96 \xi_3^2 \xi_6 - 2304 \xi_1 \xi_4 \xi_6 + 2304 \xi_1 \xi_5 \xi_6 + 768 \xi_6^2) + \\
& 96 \theta_2 (36 \xi_1^7 - 144 \xi_1^5 \xi_2 + 180 \xi_1^3 \xi_2^2 - 72 \xi_1 \xi_2^3 - 36 \xi_1^5 \xi_3 + 108 \xi_1^3 \xi_2 \xi_3 - 72 \xi_1 \xi_2^2 \xi_3 + \\
& 5 \xi_1^3 \xi_3^2 - 30 \xi_1 \xi_2 \xi_3^2 - \xi_1 \xi_3^3 - 80 \xi_1^4 \xi_4 + 144 \xi_1^2 \xi_2 \xi_4 + 72 \xi_1^2 \xi_3 \xi_4 + 48 \xi_1 \xi_4^2 + \\
& 144 \xi_1^4 \xi_5 - 144 \xi_1^2 \xi_2 \xi_5 + 136 \xi_1^2 \xi_3 \xi_5 + 48 \xi_2 \xi_3 \xi_5 + 4 \xi_3^2 \xi_5 - 480 \xi_1 \xi_4 \xi_5 - \\
& 336 \xi_1 \xi_5^2 - 288 \xi_1^3 \xi_6 + 288 \xi_1 \xi_2 \xi_6 + 16 \xi_1 \xi_3 \xi_6 + 64 \xi_4 \xi_6 - 64 \xi_5 \xi_6) + \\
& 16 \theta_3 (36 \xi_1^6 - 144 \xi_1^4 \xi_2 + 180 \xi_1^2 \xi_2^2 - 72 \xi_2^3 - 36 \xi_1^4 \xi_3 + 108 \xi_1^2 \xi_2 \xi_3 - 72 \xi_2^2 \xi_3 + \\
& 13 \xi_1^2 \xi_3^2 - 6 \xi_2 \xi_3^2 + \xi_3^3 - 80 \xi_1^3 \xi_4 + 144 \xi_1 \xi_2 \xi_4 - 24 \xi_1 \xi_3 \xi_4 + 48 \xi_4^2 + \\
& 1296 \xi_1^3 \xi_5 - 1872 \xi_1 \xi_2 \xi_5 - 72 \xi_1 \xi_3 \xi_5 - 480 \xi_4 \xi_5 - 336 \xi_5^2 + 192 \xi_1^2 \xi_6 - 48 \xi_3 \xi_6) - \\
& 512 \theta_4 (3 \xi_1^2 + 9 \xi_2 + \xi_3)];
\end{aligned} \tag{B5}$$

$$\theta_4 = \theta_1 \theta_3. \tag{B6}$$

2. Explicit form of the syzygy for the the rotation subgroup of G

The invariants $\lambda_1, \dots, \lambda_8$ of the subgroup $\mathbf{SO}_3$ of G , defined in (49), satisfy the following syzygy:

$$\begin{aligned}
\lambda_8^3 = & \frac{1}{72} (36 \lambda_1^6 - 36 \lambda_1^4 \lambda_2^2 + 9 \lambda_1^2 \lambda_2^4 - 36 \lambda_1^4 \lambda_3^2 + 18 \lambda_1^2 \lambda_2^2 \lambda_3^2 + 9 \lambda_1^2 \lambda_3^4 - \\
& 80 \lambda_1^3 \lambda_4^2 + 24 \lambda_1 \lambda_2^2 \lambda_4^2 - 16 \lambda_2^3 \lambda_4^2 + 24 \lambda_1 \lambda_3^2 \lambda_4^2 + 48 \lambda_2 \lambda_3^2 \lambda_4^2 + 48 \lambda_4^4 - \\
& 96 \lambda_2^2 \lambda_3 \lambda_4 \lambda_5 + 32 \lambda_3^3 \lambda_4 \lambda_5 - 80 \lambda_1^3 \lambda_5^2 + 24 \lambda_1 \lambda_2^2 \lambda_5^2 + 16 \lambda_2^3 \lambda_5^2 + 24 \lambda_1 \lambda_3^2 \lambda_5^2 - \\
& 48 \lambda_2 \lambda_3^2 \lambda_5^2 + 96 \lambda_4^2 \lambda_5^2 + 48 \lambda_5^4 + 96 \lambda_1^2 \lambda_2 \lambda_4 \lambda_6 + 96 \lambda_1 \lambda_2^2 \lambda_4 \lambda_6 - 48 \lambda_2^3 \lambda_4 \lambda_6 - \\
& 96 \lambda_1 \lambda_3^2 \lambda_4 \lambda_6 - 48 \lambda_2 \lambda_3^2 \lambda_4 \lambda_6 + 96 \lambda_1^2 \lambda_3 \lambda_5 \lambda_6 + 192 \lambda_1 \lambda_2 \lambda_3 \lambda_5 \lambda_6 - 48 \lambda_2^2 \lambda_3 \lambda_5 \lambda_6 - \\
& 48 \lambda_3^3 \lambda_5 \lambda_6 + 144 \lambda_1^3 \lambda_6^2 - 288 \lambda_1^2 \lambda_2 \lambda_6^2 + 72 \lambda_1 \lambda_2^2 \lambda_6^2 + 72 \lambda_1 \lambda_3^2 \lambda_6^2 - \\
& 288 \lambda_4^2 \lambda_6^2 - 288 \lambda_5^2 \lambda_6^2 + 384 \lambda_4 \lambda_6^3 - 144 \lambda_6^4 - 96 \lambda_1^2 \lambda_3 \lambda_4 \lambda_7 + 192 \lambda_1 \lambda_2 \lambda_3 \lambda_4 \lambda_7 + \\
& 48 \lambda_2^2 \lambda_3 \lambda_4 \lambda_7 + 48 \lambda_3^3 \lambda_4 \lambda_7 + 96 \lambda_1^2 \lambda_2 \lambda_5 \lambda_7 - 96 \lambda_1 \lambda_2^2 \lambda_5 \lambda_7 - 48 \lambda_2^3 \lambda_5 \lambda_7 + \\
& 96 \lambda_1 \lambda_3^2 \lambda_5 \lambda_7 - 48 \lambda_2 \lambda_3^2 \lambda_5 \lambda_7 - 576 \lambda_1^2 \lambda_3 \lambda_6 \lambda_7 + 1152 \lambda_5 \lambda_6^2 \lambda_7 + 144 \lambda_1^3 \lambda_7^2 + \\
& 288 \lambda_1^2 \lambda_2 \lambda_7^2 + 72 \lambda_1 \lambda_2^2 \lambda_7^2 + 72 \lambda_1 \lambda_3^2 \lambda_7^2 - 288 \lambda_4^2 \lambda_7^2 - 288 \lambda_5^2 \lambda_7^2 - \\
& 1152 \lambda_4 \lambda_6 \lambda_7^2 - 288 \lambda_6^2 \lambda_7^2 - 384 \lambda_5 \lambda_7^3 - 144 \lambda_7^4 - 144 \lambda_1^4 \lambda_8 + 108 \lambda_1^2 \lambda_2^2 \lambda_8 - \\
& 18 \lambda_2^4 \lambda_8 + 108 \lambda_1^2 \lambda_3^2 \lambda_8 - 36 \lambda_2^2 \lambda_3^2 \lambda_8 - 18 \lambda_3^4 \lambda_8 + 144 \lambda_1 \lambda_4^2 \lambda_8 + \\
& 144 \lambda_1 \lambda_5^2 \lambda_8 - 288 \lambda_2 \lambda_4 \lambda_6 \lambda_8 - 288 \lambda_3 \lambda_5 \lambda_6 \lambda_8 - 144 \lambda_1 \lambda_6^2 \lambda_8 + 288 \lambda_2 \lambda_6^2 \lambda_8 + \\
& 288 \lambda_3 \lambda_4 \lambda_7 \lambda_8 - 288 \lambda_2 \lambda_5 \lambda_7 \lambda_8 + 576 \lambda_3 \lambda_6 \lambda_7 \lambda_8 - 144 \lambda_1 \lambda_7^2 \lambda_8 - 288 \lambda_2 \lambda_7^2 \lambda_8 + \\
& 180 \lambda_1^2 \lambda_8^2 - 72 \lambda_2^2 \lambda_8^2 - 72 \lambda_3^2 \lambda_8^2) .
\end{aligned} \tag{B7}$$

APPENDIX C: EXPLICIT FORM OF THE $\hat{P}$ -MATRIX FOR THE GROUP G

For

$$(d_1, \dots, d_9) = (2, 4, 4, 6, 6, 8, 8, 10, 12), \tag{C1}$$

the explicit form of the $\hat{P}$ -matrix related to the integrity basis $\{p\}$ of the group G is the following

$$\begin{aligned}
P_{1,i} &= 2 d_i p_i, \quad i = 1, \dots, 9, \\
P_{2,2} &= -\frac{8}{3} (3 p_1^3 - 9 p_1 p_2 - 2 p_4 + 2 p_5), \\
P_{2,3} &= 8 p_1 p_3, \\
P_{2,4} &= 4 (2 p_1 p_4 + p_6), \\
P_{2,5} &= \frac{4}{3} (14 p_1 p_5 - 3 p_6 - 2 p_7), \\
P_{2,6} &= -\frac{2}{3} (3 p_3 p_4 - 3 p_3 p_5 - 26 p_1 p_6 + 2 p_8), \\
P_{2,7} &= -\frac{4}{3} (2 p_3 p_5 - 17 p_1 p_7 + 3 p_8), \\
P_{2,8} &= -\frac{2}{3} (2 p_3 p_6 - 3 p_3 p_7 - 32 p_1 p_8 + 3 p_9), \\
P_{2,9} &= 4 (p_3 p_8 + 5 p_1 p_9), \\
P_{3,3} &= 16 p_1 p_3, \\
P_{3,4} &= 24 p_6, \\
P_{3,5} &= 8 (p_6 + 2 p_7), \\
P_{3,6} &= 4 (p_3 p_4 + 3 p_3 p_5 + 2 p_1 p_6 + 2 p_8), \\
P_{3,7} &= 8 (2 p_3 p_5 + p_1 p_7 + p_8), \\
P_{3,8} &= 4 (2 p_3 p_6 + 3 p_3 p_7 + 4 p_1 p_8 + p_9), \\
P_{3,9} &= 24 (p_3 p_8 + p_1 p_9), \\
P_{4,4} &= 3 (6 p_1^2 - 6 p_2 - p_3) p_4, \\
P_{4,5} &= 2 (2 p_1 p_6 + p_8), \\
P_{4,6} &= \frac{1}{2} (4 p_1 p_3 p_4 + 24 p_4 p_5 + 18 p_1^2 p_6 - 18 p_2 p_6 - 3 p_3 p_6 + 2 p_9), \\
P_{4,7} &= \frac{1}{32} (-36 p_1^6 + 144 p_1^4 p_2 - 180 p_1^2 p_2^2 + 72 p_2^3 + 36 p_1^4 p_3 - 108 p_1^2 p_2 p_3 + 72 p_2^2 p_3 -
\end{aligned}$$

$$\begin{aligned}
& 9p_1^2p_3^2 + 18p_2p_3^2 + 80p_1^3p_4 - 144p_1p_2p_4 - 24p_1p_3p_4 - 48p_4^2 - 144p_1^3p_5 + \\
& 144p_1p_2p_5 - 72p_1p_3p_5 + 288p_4p_5 + 144p_5^2 - 96p_1^2p_6 + 288p_2p_6 + 112p_3p_6 + \\
& 288p_1^2p_7 - 288p_2p_7 + 32p_1p_8 + 16p_9), \\
P_{4,8} &= \frac{1}{2} (2p_3^2p_4 + 48p_4p_7 + 18p_1^2p_8 - 18p_2p_8 - 3p_3p_8 + 4p_1p_9), \\
P_{4,9} &= 3 (12p_4p_8 + 6p_1^2p_9 - 6p_2p_9 - p_3p_9), \\
P_{5,5} &= \frac{1}{3} (2p_1^2p_5 + 6p_2p_5 + 5p_3p_5 + 8p_1p_7), \\
P_{5,6} &= \frac{1}{48} (-36p_1^6 + 144p_1^4p_2 - 180p_1^2p_2^2 + 72p_2^3 + 36p_1^4p_3 - 108p_1^2p_2p_3 + 72p_2^2p_3 - \\
& 9p_1^2p_3^2 + 18p_2p_3^2 + 80p_1^3p_4 - 144p_1p_2p_4 - 24p_1p_3p_4 - 48p_4^2 - 144p_1^3p_5 + \\
& 144p_1p_2p_5 + 24p_1p_3p_5 + 480p_4p_5 + 144p_5^2 - 80p_1^2p_6 + 336p_2p_6 + 88p_3p_6 + \\
& 288p_1^2p_7 - 288p_2p_7 + 48p_3p_7 - 32p_1p_8 + 16p_9), \\
P_{5,7} &= \frac{1}{96} (-36p_1^6 + 144p_1^4p_2 - 180p_1^2p_2^2 + 72p_2^3 + 36p_1^4p_3 - 108p_1^2p_2p_3 + \\
& 72p_2^2p_3 - 9p_1^2p_3^2 + 18p_2p_3^2 + 80p_1^3p_4 - 144p_1p_2p_4 - 24p_1p_3p_4 - \\
& 48p_4^2 - 144p_1^3p_5 + 144p_1p_2p_5 + 184p_1p_3p_5 + 288p_4p_5 + 912p_5^2 - 96p_1^2p_6 + \\
& 288p_2p_6 + 48p_3p_6 + 352p_1^2p_7 - 96p_2p_7 + 160p_3p_7 - 96p_1p_8 + 16p_9), \\
P_{5,8} &= \frac{1}{6} (6p_3^2p_5 + 8p_1p_3p_6 + 96p_5p_6 + 12p_1p_3p_7 + 48p_4p_7 + 2p_1^2p_8 + 6p_2p_8 + 5p_3p_8), \\
P_{5,9} &= 2 (-12p_3p_4p_5 + p_3^2p_6 + 24p_6^2 + 2p_1p_3p_8 + 6p_4p_8), \\
P_{6,6} &= \frac{1}{12} (2p_1^2p_3p_4 + 6p_2p_3p_4 + 5p_3^2p_4 + 54p_1^2p_3p_5 - 54p_2p_3p_5 - 9p_3^2p_5 + 48p_1p_4p_5 + \\
& 24p_1p_3p_6 + 48p_4p_6 + 144p_5p_6 + 96p_4p_7 + 12p_3p_8 + 8p_1p_9), \\
P_{6,7} &= \frac{1}{96} (-36p_1^7 + 144p_1^5p_2 - 180p_1^3p_2^2 + 72p_1p_2^3 + 36p_1^5p_3 - 108p_1^3p_2p_3 + \\
& 72p_1p_2^2p_3 - 9p_1^3p_3^2 + 18p_1p_2p_3^2 + 80p_1^4p_4 - 144p_1^2p_2p_4 - 24p_1^2p_3p_4 - 48p_1p_4^2 - \\
& 144p_1^4p_5 + 144p_1^2p_2p_5 - 72p_1^2p_3p_5 + 96p_3^2p_5 + 288p_1p_4p_5 + 144p_1p_5^2 - 96p_1^3p_6 + \\
& 288p_1p_2p_6 + 176p_1p_3p_6 + 1152p_5p_6 + 288p_1^3p_7 - 288p_1p_2p_7 + 192p_1p_3p_7 + \\
& 576p_4p_7 + 576p_5p_7 - 64p_1^2p_8 + 96p_2p_8 + 80p_3p_8 + 16p_1p_9), \\
P_{6,8} &= \frac{1}{12} (8p_1p_3^2p_4 + 48p_3p_4p_5 + 12p_3^2p_6 + 96p_6^2 + 54p_1^2p_3p_7 - 54p_2p_3p_7 - 9p_3^2p_7 + \\
& 96p_1p_4p_7 + 24p_1p_3p_8 + 72p_4p_8 + 216p_5p_8 + 2p_1^2p_9 + 6p_2p_9 + 5p_3p_9), \\
P_{6,9} &= \frac{1}{2} (2p_3^3p_4 + 24p_3p_4p_6 + 18p_1^2p_3p_8 - 18p_2p_3p_8 - 3p_3^2p_8 + 24p_1p_4p_8 + 4p_1p_3p_9 + \\
& 12p_4p_9 + 60p_5p_9), \\
P_{7,7} &= \frac{1}{3} (2p_1^2p_3p_5 + 6p_2p_3p_5 + 5p_3^2p_5 + 12p_1p_5^2 + 24p_5p_6 + 8p_1p_3p_7 + 48p_5p_7), \\
P_{7,8} &= \frac{1}{6} (12p_1p_3^2p_5 + 36p_3p_4p_5 - 36p_3p_5^2 + 2p_1^2p_3p_6 + 6p_2p_3p_6 + 5p_3^2p_6 + 48p_1p_5p_6 + \\
& 24p_6^2 + 6p_3^2p_7 + 72p_7^2 + 8p_1p_3p_8 + 120p_5p_8), \\
P_{7,9} &= -12p_1p_3p_4p_5 + 4p_1p_3^2p_6 + 12p_3p_4p_6 - 12p_3p_5p_6 + 24p_1p_6^2 + 2p_3^2p_8 + 24p_7p_8 + \\
& 24p_5p_9, \\
P_{8,8} &= \frac{1}{12} (2p_1^2p_3^2p_4 + 6p_2p_3^2p_4 + 5p_3^3p_4 + 54p_1^2p_3^2p_5 - 54p_2p_3^2p_5 - 9p_3^3p_5 + \\
& 192p_1p_3p_4p_5 + 24p_1p_3^2p_6 + 96p_3p_4p_6 - 288p_3p_5p_6 + 12p_3^2p_8 + 576p_7p_8 + \\
& 8p_1p_3p_9 + 192p_5p_9), \\
P_{8,9} &= \frac{1}{2} (4p_1p_3^3p_4 + 12p_3^2p_4^2 + 18p_1^2p_3^2p_6 - 18p_2p_3^2p_6 - 3p_3^3p_6 + 48p_1p_3p_4p_6 -
\end{aligned}$$

$$\begin{aligned}
& 84 p_3 p_6^2 - 24 p_3 p_4 p_8 + 84 p_8^2 + 2 p_3^2 p_9 + 48 p_6 p_9 + 84 p_7 p_9), \\
P_{9,9} = & 3 (6 p_1^2 p_3^3 p_4 - 6 p_2 p_3^3 p_4 - p_3^4 p_4 + 12 p_1 p_3^2 p_4^2 + 288 p_1 p_3^2 p_4 p_5 - 1152 p_3 p_4 p_5^2 - \\
& 24 p_3^2 p_4 p_6 - 288 p_1 p_3 p_6^2 + 1152 p_5 p_6^2 + 144 p_1^2 p_3 p_4 p_7 - 432 p_2 p_3 p_4 p_7 - 72 p_3^2 p_4 p_7 + \\
& 1152 p_4 p_7^2 + 108 p_1^6 p_8 - 432 p_1^4 p_2 p_8 + 540 p_1^2 p_2^2 p_8 - 216 p_2^3 p_8 - 108 p_1^4 p_3 p_8 + \\
& 324 p_1^2 p_2 p_3 p_8 - 216 p_2^2 p_3 p_8 + 27 p_1^2 p_3^2 p_8 - 54 p_2 p_3^2 p_8 - 240 p_1^3 p_4 p_8 + \\
& 432 p_1 p_2 p_4 p_8 + 72 p_1 p_3 p_4 p_8 + 144 p_4^2 p_8 + 432 p_1^3 p_5 p_8 - 432 p_1 p_2 p_5 p_8 + \\
& 216 p_1 p_3 p_5 p_8 - 864 p_4 p_5 p_8 - 432 p_5^2 p_8 - 864 p_1^2 p_7 p_8 + 864 p_2 p_7 p_8 + 144 p_1^2 p_5 p_9 - \\
& 432 p_2 p_5 p_9 - 72 p_3 p_5 p_9 + 288 p_1 p_7 p_9).
\end{aligned}$$

APPENDIX D: EXPLICIT FORM OF THE GENERATORS OF THE ISOTROPY SUBGROUPS OF $\mathbf{SO}_3$

In this Appendix are collected the definitions of the groups and of the elements and the generators of the subgroups of $\mathbf{SO}_3$, involved in the body of the paper, in the tables and in the figures.

a. Generators or group elements

$$\begin{aligned}
C_{2x} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, & C_{2z} &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & C_{3z} &= \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} & 0 \\ -\frac{\sqrt{3}}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\
C_{4z} &= \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & C_{4x} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, & C_{2a} &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \\
C_{3\delta} &= \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, & R_z(\phi) &= \begin{pmatrix} \cos(\phi) & \sin(\phi) & 0 \\ -\sin(\phi) & \cos(\phi) & 0 \\ 0 & 0 & 1 \end{pmatrix}, & R_x(\phi) &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\phi) & \sin(\phi) \\ 0 & -\sin(\phi) & \cos(\phi) \end{pmatrix}.
\end{aligned} \tag{D1}$$

b. Groups

By $\mathbf{SO}_3$ and $\mathbf{O}_3$ we denote the proper and, respectively, the complete 3D rotation group and, by $\mathbf{O}_2^\alpha$, $\alpha = x, z$ the following $\mathbf{O}_2$ subgroups of $\mathbf{SO}_3$: $\mathbf{O}_2^x = \{R_x(\phi)\}_{0 \leq \phi < 2\pi} \cup \{C_{2z} R_x(\phi)\}_{0 \leq \phi < 2\pi}$ and $\mathbf{O}_2^z = \{R_z(\phi)\}_{0 \leq \phi < 2\pi} \cup \{C_{2x} R_z(\phi)\}_{0 \leq \phi < 2\pi}$.

-
- [1] P. W. Anderson, P. Morel Phys. Rev. **123**, 1911 (1961).
[2] D. D. Osheroff, R.C. Richardson, D.M. Lee, Phys. Rev. Lett. **28**, 885 (1972).
[3] M. Sigrist, K. Ueda, Rev. Mod. Phys. **63**, 239 (1991).
[4] D. J. van Harlingen, Rev. Mod. Phys. **67**, 515 (1995).
[5] H. Ghosh, J. Phys. Condens. Matter **11**, L371 (1999).
[6] J. Orenstein, Nature **401**, 333 (1999).
[7] J.F. Annet, Physica **C 317-318**, 1 (1999).
[8] Yu. M. Gufan, G. M. Vereshkov, P. Tolédano, B. Mettout, R. Bouzerar and V. Lorman, Phys. Rev. **B51**, 9219 and 9228 (1995).
[9] K. Nakamura, A. Yu. Gufan, Yu. M. Gufan and E. G. Rudashevskii, Crystallography Reports **44**, 469 and 603 (1999).
[10] Yu. M. Gufan, I. G. Levchenko and E. G. Rudashevskii, Physics of the Solid State **41**, 1422 (1999) and references therein.
[11] N. D. Mermin, Phys. Rev. A **9**, 868 (1974).
[12] A. M. J. Schakel, F. A. Bias, J. Phys.: Condens. Matter **1**, 1743(1989).
[13] H. W. Capel, A. M. J. Schakel, Physica A **160**, 409 (1989).
[14] M. Abud and G. Sartori, Phys. Lett. **B 104**, 147 (1981) and Ann. Phys. **150**, 307 (1983).
[15] Yu. M. Gufan, Sov. Phys. Solid State **13**, 175 (1971).
[16] G. Sartori, Supplmento del Nuovo Cimento **14**, 1 (1991).
[17] C. Procesi and G. W. Schwarz, Inv. Math. **81**, 539 (1985).
[18] E. B. Vinberg and V. L. Popov, *Invariant Theory*, in Encyclopaedia of Mathematical Sciences, Vol. **55** (Springer-Verlag, Berlin, 1994), pp. 123-284.
[19] V. L. Popov, *Groups, Generators, Syzygies and Orbits in Invariant Theory*, Trans. Math. Monog. Vol. **100**, (A.M.S., Providence, 1992).
[20] R. Stanley, Bull. Am. Math. Soc. New. Ser. **1**, 475 (1979).
[21] G. Sartori and G. Valente, in preparation.
[22] J. F. Cornwell, *Group Theory in Physics*, (Academic, London, 1984),Vol. I.
[23] G. Sartori and G. Valente, J. Phys. A **29**, 193 (1996).
[24] G. Sartori and V. Talamini, J. Math. Phys. **39**, 2367 (1998).
[25] R. J. Walker, *Algebraic Curves*, (Dover, New York, 1950) (Theorem. 5.1 p. 67).

TABLE I. Relations defining 1D geometrical strata, $\widehat{S}^{(1,A)} = \widehat{W}^{(1,A)}$ in orbit space. For $2 \leq i \leq 9$, $\tilde{p}_i = p_i/(p_1^{d_i/2})$, d_i denotes the degree of the polynomial $p_i(x)$ in the order parameter x and the values, $p^{(1,A)}$, taken on by $\tilde{p}$ on $\widehat{W}^{(1,A)}$ are listed in the columns from 2 to 9.

A	$\tilde{p}_2$	$\tilde{p}_3$	$\tilde{p}_4$	$\tilde{p}_5$	$\tilde{p}_6$	$\tilde{p}_7$	$\tilde{p}_8$	$\tilde{p}_9$
1	1	0	0	0	0	0	0	0
2	1/2	1	1/6	1/6	1/6	1/6	1/6	1/6
3	1/2	1	0	0	0	0	0	0
4	1/3	0	1/3	0	0	0	0	0
5	1/2	0	0	0	0	0	0	0

TABLE II. Parametric equations defining the geometrical strata, $\widehat{W}^{(2,A)}$ (A is listed in the head row) in orbit space. For $2 \leq i \leq 9$, $\tilde{p}_i = p_i/(p_1^{d_i/2})$ and d_i denotes the degree of the polynomial $p_i(x)$, in the order parameter x , and $\epsilon = \pm 1$.

$\tilde{p} \backslash A$	1_ϵ	2	3_ϵ	4	5
$\tilde{p}_2$	$(2 + \xi^2)/6$	$1/2$	$(2 + \xi^2)/6$	$1/2$	ξ
$\tilde{p}_3$	0	ξ	ξ^2	1	$2 - 2\xi$
$\tilde{p}_4$	$(2 - \xi)^2(1 + \xi)/12$	0	$(2 - \xi)^2(1 + \xi)/12$	ξ	0
$\tilde{p}_5$	0	0	$\xi^2(1 + \xi)/12$	ξ	0
$\tilde{p}_6$	0	0	$(2 - \xi)(1 + \xi)\xi^2/12$	ξ	0
$\tilde{p}_7$	0	0	$(1 + \xi)\xi^3/12$	ξ	0
$\tilde{p}_8$	0	0	$(2 - \xi)(1 + \xi)\xi^3/12$	ξ	0
$\tilde{p}_9$	0	0	$(2 - \xi)^2(1 + \xi)\xi^3/12$	ξ	0
ξ range	$]0, \frac{1+3\epsilon}{2}[$	$]0, 1[$	$]0, \epsilon[$	$]0, \frac{1}{6}[$	$] \frac{1}{2}, 1[$

TABLE III. Possible symmetry strata, $S^{(d,r)}$ for D-wave driven pairing in isotropic space (d denotes the dimension of the stratum and r is an enumeration index). From left to right, the columns refer to the phase reference numbers according to our $((d, r))$ and ref.[12] ($\mathcal{N}$) classification, the order ($|H|$) and the the residual symmetry group, H , the complex coordinates ($z_j = x_j + i x_{5+j}$, $j = 1, \dots, 5$) of the H -invariant superconducting vector order parameter and corresponding values of the partial wave amplitudes ($D_2, \dots, D_{-2}$). In columns five and six, the t 's are real parameters, while the v 's are complex ones. The definitions of the group elements are recalled in Appendix D.

(d, r)	$\mathcal{N}$	$ H $	H	$(z_1, \dots, z_5)$	$\sqrt{2}(D_2, D_1, D_0, D_{-1}, D_{-2})$
(1,1)	II	∞	$\langle C_{2x} \mathcal{T} \rangle \times \{R_z(\phi) U_1(2\phi)\}_\phi$	$(it, -t, 0, 0, 0)$	$(2t, 0, 0, 0, 0)$
(1,2)	$VIII$	∞	$\mathbf{O}_2^z \otimes \langle \mathcal{T} \rangle$	$(0, 0, 0, 0, t)$	$(0, 0, t\sqrt{2}, 0, 0)$
(1,3)	IX, X	16	$\langle C_{2x}, \mathcal{T}, C_{4z} U_1(\pi) \rangle$	$(0, t, 0, 0, 0)$	$(-t, 0, 0, 0, -t)$
(1,4)	XI	24	$\langle C_{2x}, C_{2a} \mathcal{T}, C_{3\delta} U_1(4\pi/3) \rangle$	$(0, -it, 0, 0, t)$	$(it, 0, t\sqrt{2}, 0, it)$
(1,5)	I	∞	$\langle C_{2x} \mathcal{T} \rangle \times \{R_z(\phi) U_1(-\phi)\}_\phi$	$(0, 0, t, -it, 0)$	$(0, 0, 0, 2it, 0)$
(2,1)	V	6	$\langle C_{2x} \mathcal{T}, C_{3z} U_1(4\pi/3) \rangle$	$(it_1, -t_1, t_2, -it_2, 0)$	$(2t_1, 0, 0, 2it_2, 0)$
(2,2)	IV	4	$\langle C_{2x} \mathcal{T}, C_{2z} U_1(\pi) \rangle$	$(0, 0, t_1, it_2, 0)$	$(0, it_1 + it_2, 0, it_1 - it_2)$
(2,3)	$--$	8	$\langle C_{2x}, C_{4z} \mathcal{T} \rangle$	$(0, it_1, 0, 0, t_2)$	$(-it_1, 0, t_2\sqrt{2}, 0, -it_1)$
(2,4)	VII	8	$\langle C_{2x}, C_{2z}, \mathcal{T} \rangle$	$(0, t_1, 0, 0, t_2)$	$(-t_1, 0, t_2\sqrt{2}, 0, -t_1)$
(2,5)	VI	8	$\langle C_{2x} \mathcal{T}, C_{4z} U_1(\pi) \rangle$	$(it_1, t_2, 0, 0, 0)$	$(t_1 - t_2, 0, 0, 0, -t_1 - t_2)$
(3,1)	III	4	$\langle C_{2z}, C_{2x} \mathcal{T} \rangle$	$(it_1, t_2, 0, 0, t_3)$	$(t_1 - t_2, 0, t_3\sqrt{2}, 0, -t_1 - t_2)$
(3,2)	$--$	4	$\langle C_{2x}, C_{2z} \rangle$	$(0, v_1, 0, 0, v_2)$	$(-v_1, 0, v_2\sqrt{2}, 0, -v_1)$
(4,1)	$--$	2	$\langle C_{2x} \mathcal{T} \rangle$	$(it_1, t_2, t_3, it_4, t_5)$	$(t_1 - t_2, it_3 + it_4, t_5\sqrt{2}, it_3 - it_4, -t_1 - t_2)$
(4,2)	$--$	2	$\langle C_{2z} \rangle$	$(v_1, v_2, 0, 0, v_3)$	$(-iv_1 - v_2, 0, v_3\sqrt{2}, 0, iv_1 - v_2)$
(6,1)	$--$	1	$\{1\}$	$(v_1, v_2, v_3, v_4, v_5)$	$(-iv_1 - v_2, iv_3 + v_4, v_5\sqrt{2}, iv_3 - v_4, iv_1 - v_2)$

TABLE IV. Strata of dimensions ≥ 3 , bordering each stratum $\widehat{S}^{(d,r)}$ (d denotes the dimension of the stratum and r is an enumeration index). From left to right, the columns refer to the enumeration numbers, (d, r) , of the lower bordering stratum, $\widehat{S}^{(d,r)}$, its orbit-type, $[G^{(d,r)}]$, the maximal conjugacy classes of subgroups of $[G^{(d,r)}]$, $[H]_M$, the order, $|H_M|$, of H_M and the indices, (d', r') , of the corresponding strata, if any. The definitions of the group elements are recalled in Appendix D.

(d, r)	$[G^{(d,r)}]$	$[H]_M$	$ H_M $	(d', r')
(2,1)	$[\langle C_{2x}T, C_{3z}U_1(4\pi/3) \rangle]$	$[\langle C_{3z}U_1(4\pi/3) \rangle]$	3	–
		$[\langle C_{2x}T \rangle]$	2	(4,1)
(2,2)	$[\langle C_{2x}T, C_{2z}U_1(\pi) \rangle]$	$[\langle C_{2x}T \rangle]$	2	(4,1)
		$[\langle C_{2z}U_1(\pi) \rangle]$	2	–
(2,3)	$[\langle C_{2z}, C_{4x}T \rangle]$	$[\langle C_{2z}, C_{2x}T \rangle]$	4	(3,1)
		$[\langle C_{2x}, C_{2z} \rangle]$	4	(3,2)
		$[\langle C_{4x}T \rangle]$	4	–
(2,4)	$[\langle C_{2x}, C_{2z}, T \rangle]$	$[\langle C_{2x}, C_{2z} \rangle]$	4	(3,2)
		$[\langle C_{2z}, C_{2x}T \rangle]$	4	(3,1)
(2,5)	$[\langle C_{2x}T, C_{4z}U_1(\pi) \rangle]$	$[\langle C_{2z}, C_{2x}T \rangle]$	4	(3,1)
		$[\langle C_{4z}U_1(\pi) \rangle]$	4	–
(3,1)	$[\langle C_{2z}, C_{2x}T \rangle]$	$[\langle C_{2x}T \rangle]$	2	(4,1)
		$[\langle C_{2z} \rangle]$	2	(4,2)
(3,2)	$[\langle C_{2x}, C_{2z} \rangle]$	$[\langle C_{2z} \rangle]$	2	(4,2)
(4,1)	$[\langle C_{2x}T \rangle]$	$[1]$	1	(6,1)
(4,2)	$[\langle C_{2z} \rangle]$	$[1]$	1	(6,1)

TABLE V. Absolute minimum, $\widehat{\Phi}_{\min}^{(4)} = -\alpha_1^2/(2\delta)$, of a general, bounded below, G -invariant 4-th degree polynomial, $\widehat{\Phi}^{(4)}(\alpha, p) = \alpha_0 p_1^2/2 + \sum_{j=1}^3 \alpha_j p_j$, $\alpha_1 < 0$, and hosting strata, $S^{(d,r)}$, as functions of the coefficients α . The denomination of the strata is the same as in Table III.

α range	δ	(d, r)
$\mathcal{R}_1 : \text{Max}(0, -3\alpha_0/2, -6\alpha_3) < \alpha_2$	$\alpha_0 + 2\alpha_2/3$	(1,4)
$\mathcal{R}_2 : -6\alpha_3 > \alpha_2 > \text{Max}(-\alpha_0 - 2\alpha_3, 2\alpha_3)$	$\alpha_0 + \alpha_2 + 2\alpha_3$	(1,2), (1,3), (2,4)
$\mathcal{R}_3 : -\alpha_0/2 < \alpha_2 < \text{Min}(0, 2\alpha_3)$	$\alpha_0 + 2\alpha_2$	(1,1)
$\mathcal{R}_{13} : 0 = \alpha_2 < \text{Min}(\alpha_0, \alpha_3)$	α_0	(1,1), (1,4), (1,5), (2,1), (3,1), (4,1)
$\mathcal{R}_{12} : \text{Max}(-3\alpha_0/2, 0) < \alpha_2 = -6\alpha_3$	$\alpha_0 - 4\alpha_3$	(1,2), (1,3), (1,4), (2,3), (2,4), (3,2)
$\mathcal{R}_{23} : -\alpha_0/2 < \alpha_2 = 2\alpha_3 < 0$	$\alpha_0 + 4\alpha_3$	(1,1), (1,2), (1,3), (2,4), (2,5)
$\mathcal{R}_{123} : \alpha_2 = \alpha_3 = 0 < \alpha_0$	α_0	all, except (0,1)

FIG. 1. Possible phase transitions between bordering strata, connected, in the figure, by continuous sequences of one or more arrows. The group elements are defined in Appendix D.




FIG. 2. Localization of the absolute minimum of a fourth degree G -invariant polynomial, $\widehat{\Phi}^{(4)}(\alpha, p) = \alpha_0 p_1^2/2 + \sum_{j=1}^3 \alpha_j p_j$, $\alpha_1 < 0$, as a function of its coefficients. For values of $(\alpha_0, \alpha_2, \alpha_3)$ in $\mathcal{R}_1$ or in $\mathcal{R}_2$ or in $\mathcal{R}_3$, the absolute minimum lies, respectively, in the strata $S^{(1,4)}$ or $\{S^{(1,2)}, S^{(1,3)}, S^{(2,4)}\}$ (degenerate minimum) or $S^{(1,1)}$. For particular values of the α 's, see Table IV.

